

# Document made available under the Patent Cooperation Treaty (PCT)

International application number: PCT/EP05/001432

International filing date: 07 February 2005 (07.02.2005)

Document type: Certified copy of priority document

Document details: Country/Office: GB  
Number: 0402809.8  
Filing date: 09 February 2004 (09.02.2004)

Date of receipt at the International Bureau: 29 April 2005 (29.04.2005)

Remark: Priority document submitted or transmitted to the International Bureau in compliance with Rule 17.1(a) or (b)



World Intellectual Property Organization (WIPO) - Geneva, Switzerland  
Organisation Mondiale de la Propriété Intellectuelle (OMPI) - Genève, Suisse



INVESTOR IN PEOPLE

EPO5 / 1432


The Patent Office  
Concept House  
Cardiff Road  
Newport  
South Wales  
NP10 8QQ

I, the undersigned, being an officer duly authorised in accordance with Section 74(1) and (4) of the Deregulation & Contracting Out Act 1994, to sign and issue certificates on behalf of the Comptroller-General, hereby certify that annexed hereto is a true copy of the documents as originally filed in connection with the patent application identified therein.

In accordance with the Patents (Companies Re-registration) Rules 1982, if a company named in this certificate and any accompanying documents has re-registered under the Companies Act 1980 with the same name as that with which it was registered immediately before re-registration save for the substitution as, or inclusion as, the last part of the name of the words "public limited company" or their equivalents in Welsh, references to the name of the company in this certificate and any accompanying documents shall be treated as references to the name with which it is so re-registered.

In accordance with the rules, the words "public limited company" may be replaced by p.l.c., plc, P.L.C. or PLC.

Re-registration under the Companies Act does not constitute a new legal entity but merely subjects the company to certain additional company law rules.



Signed

Dated 7 April 2005



2000

Patents Act 1977  
(Rule 16)


**The  
Patent  
Office**

10FEB04 E8718 01-1 002025  
P01/7700 0.00 0402809.8 NONE

22-04-05  
**1/77**

**Request for grant of a patent**

(See the notes on the back of this form. You can also get an explanatory leaflet from the Patent Office to help you fill in this form)



**The Patent Office**

Cardiff Road  
Newport  
South Wales  
NP10 8QQ

1. Your Reference	SAL/PB60713P		
2. Patent application number (The Patent office will fill in this part)	0402809.8		09 FEB 2004
3. Full name, address and postcode of the or of each applicant (underline all surnames)	GLAXO GROUP LIMITED GLAXO WELLCOME HOUSE BERKELEY AVENUE GREENFORD MIDDLESEX UB6 ONN GB		
Patents ADP number (if you know it)			
If the applicant is a corporate body, give the country/state of its corporation	GB		
4 Title of the invention	CHEMICAL COMPOUNDS		
5 Name of your agent (if you have one)	STEPHANIE A. LEAROYD		
"Address for service" in the United Kingdom to which all correspondence should be sent (including the postcode)	GLAXOSMITHKLINE CORPORATE INTELLECTUAL PROPERTY 980 GREAT WEST ROAD BRENTFORD, MIDDLESEX TW8 9GS, GB		
Patents ADP number (if you know it)	8072555004		
6. Priority: Complete this section if you are declaring priority from one or more earlier patent applications, filed in the last 12 months	Country	Priority application number (if you know it)	Date of Filing (day / month / year)
7. Divisionals: etc Complete this section only if this application is a divisional application or resulted from an entitlement dispute (see note f)	Number of earlier UK application		Date of filing (day / month / year)
8. Is a Patents Form 7/77 (Statement of inventorship and of right to grant of a patent) required in support of this request?	YES		
Answer YES if:			
a) any applicant named in part 3 is not an inventor, or			
b) there is an inventor who is not named as an applicant, or			
c) any named applicant is a corporate body			
Otherwise answer NO See note (d)			

473587003

**Patents Form 1/77**

9. Accompanying documents: A patent application must include a description of the invention. Not counting duplicates, please enter the number of pages of each item accompanying this form:

Continuation sheet of this form

Description	42 <i>DL</i>
Claim(s)	7
Abstract	-
Drawing(s)	-

10. If you are also filing any of the following, state how many against each item

Priority Documents

Translations of priority documents

Statement of inventorship and right to grant of a patent (*Patents Form 7/77*)

Request for preliminary examination and search (*Patent Form 9/77*)

Request for substantive examination (*Patent Form 10/77*)

Any other documents(*please specify*)

11. I/We request the grant of a patent on the basis of this application

STEPHANIE A. LEAROYD

Signature(s)

Date: 9 FEBRUARY 2004

12. Name and daytime telephone number and e-mail address, **KATHERINE Y EVANS** **01 438 768611**  
if any, of person to contact in the United Kingdom

**Warning**

After an application for a patent has been filed, the Comptroller of the Patent Office will consider whether publication or communication of the invention should be prohibited or restricted under Section 22 of the Patents Act 1977. You will be informed if it is necessary to prohibit or restrict your invention in this way. Furthermore, if you live in the United Kingdom, Section 23 of the Patents Act 1977 stops you from applying for a patent abroad without first getting written permission from the Patent Office unless an application has been filed at least 6 weeks beforehand in the United Kingdom for a patent for the same invention and either no direction prohibiting publication or communication has been given, or any such direction has been revoked

**Notes**

- If you need help to fill in this form or you have any questions, please contact the Patent Office on 08459 500505.
- Write your answers in capital letters using black ink or you may type them.
- If there is not enough space for all the relevant details on any part of this form, please continue on a separate sheet of paper and write "see continuation sheet" in the relevant part(s). Any continuation sheet should be attached to this form
- If you have answered "Yes" in part 8, a Patents Form 7/77 will need to be filed.
- Once you have filled in the form you must remember to sign and date it.
- Part 7 should only be completed when a divisional application is being made under section 15(4), or when an application is being made under section 8(3), 12(6) or 37(4) following an entitlement dispute. By completing part 7 you are requesting that this application takes the same filing date as an earlier UK application. If you want the new application to have the same priority date(s) as the earlier UK application, you should also complete part 6 with the priority details.

## CHEMICAL COMPOUNDS

### BACKGROUND OF THE INVENTION

The present invention relates to benzimidazole derivatives, compositions and medicaments containing the same, as well as processes for the preparation and use of such compounds, compositions and medicaments. Such benzimidazole derivatives are useful in the treatment of diseases associated with inappropriate I-kappa-B kinase-3 (IKK3) (also known as I-kappa-B kinase epsilon (IKK $\epsilon$ ) or inducible I-kappa B kinase (IKKi)) activity, including in particular in the treatment and prevention of disease states mediated by IKK3 mechanisms including inflammatory and tissue repair disorders, particularly rheumatoid arthritis, inflammatory bowel disease, asthma and COPD (chronic obstructive pulmonary disease); osteoarthritis, osteoporosis and fibrotic diseases; dermatosis, including psoriasis, atopic dermatitis and ultraviolet radiation (UV)-induced skin damage; autoimmune diseases including systemic lupus erythematosus, multiple sclerosis, psoriatic arthritis, ankylosing spondylitis, tissue and organ rejection, Alzheimer's disease, stroke, atherosclerosis, restenosis, diabetes, glomerulonephritis, cancer, including Hodgkins disease, cachexia, inflammation associated with infection and certain viral infections, including acquired immune deficiency syndrome (AIDS), adult respiratory distress syndrome, and Ataxia Telangiectasia.

An important large family of enzymes is the protein kinase enzyme family. Currently, there are about 500 different known protein kinases. However, because three to four percent of the human genome is a code for the formation of protein kinases, there may be many thousands of distinct and separate kinases in the human body. Protein kinases serve to catalyze the phosphorylation of an amino acid side chain in various proteins by the transfer of the  $\gamma$ -phosphate of the ATP-Mg<sup>2+</sup> complex to said amino acid side chain. These enzymes control the majority of the signaling processes inside cells, thereby governing cell function, growth, differentiation and destruction (apoptosis) through reversible phosphorylation of the hydroxyl groups of serine, threonine and tyrosine residues in proteins. Studies have shown that protein kinases are key regulators of many cell functions, including signal transduction, transcriptional regulation, cell motility, and cell division. Several oncogenes have also been shown to encode protein kinases, suggesting that kinases play a role in oncogenesis. These processes are highly regulated, often by complex intermeshed pathways where each kinase will itself be regulated by one or more kinases. Consequently, aberrant or inappropriate protein kinase activity can contribute to the rise of disease states associated with such aberrant kinase activity. Due to their physiological relevance, variety and ubiquitousness, protein kinases have become one of the most important and widely studied family of enzymes in biochemical and medical research.


The protein kinase family of enzymes is typically classified into two main subfamilies: Protein Tyrosine Kinases and Protein Serine/Threonine Kinases, based on the amino acid

residue they phosphorylate. The serine/threonine kinases (PSTK), includes cyclic AMP- and cyclic GMP-dependent protein kinases, calcium and phospholipid dependent protein kinase, calcium- and calmodulin-dependent protein kinases, casein kinases, cell division cycle protein kinases and others. These kinases are usually cytoplasmic or associated with the particulate fractions of cells, possibly by anchoring proteins. Aberrant protein serine/threonine kinase activity has been implicated or is suspected in a number of pathologies such as rheumatoid arthritis, psoriasis, septic shock, bone loss, many cancers and other proliferative diseases. Accordingly, serine/threonine kinases and the signal transduction pathways which they are part of are important targets for drug design. The tyrosine kinases phosphorylate tyrosine residues. Tyrosine kinases play an equally important role in cell regulation. These kinases include several receptors for molecules such as growth factors and hormones, including epidermal growth factor receptor, insulin receptor, platelet derived growth factor receptor and others. Studies have indicated that many tyrosine kinases are transmembrane proteins with their receptor domains located on the outside of the cell and their kinase domains on the inside. Much work is also under progress to identify modulators of tyrosine kinases as well.

The present inventors have discovered novel benzimidazole derivatives which are inhibitors of kinase activity, in particular inappropriate IKK3 activity. Such derivatives are therefore useful in the treatment of disorders associated with inappropriate kinase activity, in particular inappropriate IKK3 activity in particular in the treatment and prevention of disease states mediated by IKK3 mechanisms including inflammatory and tissue repair disorders, particularly rheumatoid arthritis, inflammatory bowel disease, asthma and COPD (chronic obstructive pulmonary disease); osteoarthritis, osteoporosis and fibrotic diseases; dermatosis, including psoriasis, atopic dermatitis and ultraviolet radiation (UV)-induced skin damage; autoimmune diseases including systemic lupus erythematosus, multiple sclerosis, psoriatic arthritis, ankylosing spondylitis, tissue and organ rejection, Alzheimer's disease, stroke, atherosclerosis, restenosis, diabetes, glomerulonephritis, cancer, including Hodgkins disease, cachexia, inflammation associated with infection and certain viral infections, including acquired immune deficiency syndrome (AIDS), adult respiratory distress syndrome, and Ataxia Telangiectasia.

#### **BRIEF SUMMARY OF THE INVENTION**

In one aspect of the present invention, there is provided a compound of Formula (I):



wherein

n is 0, 1, 2, 3, or 4;

Each R¹ which may be the same or different, independently represents H, halogen or a group (X)<sub>a</sub>(Y)<sub>b</sub>Z;

X represents -O- or -CONH-;

a is 0 or 1;

Y represents -C<sub>1-6</sub> alkylene-

b is 0 or 1;

Z represents hydroxy, C<sub>1-6</sub> alkyl, C<sub>1-6</sub> haloalkyl, C<sub>5-7</sub> heterocyclyl, C<sub>1-6</sub> alkoxyalkyl, C<sub>1-6</sub> haloalkoxyalkyl;

R² represents a group -(X¹)<sub>c</sub>(Y¹)<sub>d</sub>Z¹

Wherein X¹ represents -C<sub>1-12</sub> alkylene-;

c is 0 or 1;

Y¹ represents -O-;

d is 0 or 1;

Z¹ represents H, aryl or heteroaryl each of which contains 5 - 14 ring atoms, C<sub>5-7</sub>


heterocyclyl, C<sub>5-7</sub> cycloalkyl, C<sub>5-7</sub> cycloalkenyl, each of which may be optionally substituted by one or more substituents independently selected from C<sub>1-6</sub> alkyl, C<sub>1-6</sub> haloalkyl, halogen, C<sub>1-6</sub> alkoxy, C<sub>1-6</sub> haloalkoxy, SO<sub>2</sub>R³, C<sub>1-6</sub> hydroxyalkyl.

R³ represents H or C<sub>1-6</sub> alkyl;

or pharmaceutically acceptable salts, solvates or physiologically functional derivatives


thereof with the proviso that the compound is not:

5-(1*H*-benzimidazol-1-yl)-3-[(2-methylphenyl)oxy]thiophene-2-carbonitrile




5-(5,6-bismethoxy-1*H*-benzimidazol-1-yl)-3-[(2-trifluoromethyl)phenyl]oxy}thiophene-2-carbonitrile





In a second aspect of the present invention, there is provided a compound of Formula (I), or a salt, solvate, or a physiologically functional derivative thereof for use in therapy.

- 5 In a third aspect of the present invention, there is provided a pharmaceutical composition comprising a therapeutically effective amount of a compound of Formula (I) or a salt, solvate, or a physiologically functional derivative thereof and one or more of pharmaceutically acceptable carriers, diluents and excipients.
- 10 In a fourth aspect of the present invention, there is provided a method of treating a disorder in a mammal, said disorder being mediated by inappropriate IKK3 activity, comprising administering to said mammal a therapeutically effective amount of a compound of Formula (Ia)



15 wherein

n is 0, 1, 2, 3, or 4;

Each R¹ which may be the same or different, independently represents H, halogen or a group (X)<sub>a</sub>(Y)<sub>b</sub>Z;

20 X represents -O- or -CONH-;

a is 0 or 1;

Y represents -C<sub>1-6</sub> alkylene-

b is 0 or 1;

25 Z represents hydroxy, C<sub>1-6</sub> alkyl, C<sub>1-6</sub> haloalkyl, C<sub>5-7</sub> heterocyclyl, C<sub>1-6</sub> alkoxyalkyl, C<sub>1-6</sub> haloalkoxyalkyl;

R² represents a group -(X¹)<sub>c</sub>(Y¹)<sub>d</sub>Z¹;

Wherein X¹ represents -C<sub>1-6</sub> alkylene-;

c is 0 or 1;


30 Y¹ represents -O-;

d is 0 or 1;

$Z^1$  represents H, aryl or heteroaryl each of which contains 5 - 14 ring members,  $C_{5-7}$  heterocyclyl,  $C_{5-7}$  cycloalkyl,  $C_{5-7}$  cycloalkenyl, each of which may be optionally substituted by one or more substituents independently selected from  $C_{1-6}$  alkyl,  $C_{1-6}$  haloalkyl, halogen,  $C_{1-6}$  alkoxy,  $C_{1-6}$  haloalkoxy,  $SO_2R^3$ ,  $C_{1-6}$  hydroxyalkyl.

- 5  $R^3$  represents H or  $C_{1-6}$  alkyl;  
or pharmaceutically acceptable salts, solvates or physiologically functional derivatives thereof.

10 In a fifth aspect of the present invention, there is provided the use of a compound of Formula (I) (a):



wherein

$n$  is 0, 1, 2, 3, or 4;

- 15 Each  $R^1$  which may be the same or different, independently represents H, halogen or a group  $(X)_a(Y)_bZ$ ;

$X$  represents  $-O-$  or  $-CONH-$ ;

$a$  is 0 or 1;

$Y$  represents  $-C_{1-6}$  alkylene-

- 20  $b$  is 0 or 1;

$Z$  represents hydroxy,  $C_{1-6}$  alkyl,  $C_{1-6}$  haloalkyl,  $C_{5-7}$  heterocyclyl,  $C_{1-6}$  alkoxyalkyl,  $C_{1-6}$  haloalkoxyalkyl;

$R^2$  represents a group  $-(X^1)_c(Y^1)_dZ^1$ ;

25

Wherein  $X^1$  represents  $-C_{1-6}$  alkylene-;

$c$  is 0 or 1;

$Y^1$  represents  $-O-$ ;

$d$  is 0 or 1;

30

$Z^1$  represents H, aryl or heteroaryl, each of which contains 5 - 14 ring members,  $C_{5-7}$  heterocyclyl,  $C_{5-7}$  cycloalkyl,  $C_{5-7}$  cycloalkenyl, each of which may be optionally substituted by one or more substituents independently selected from  $C_{1-6}$  alkyl,  $C_{1-6}$  haloalkyl, halogen,  $C_{1-6}$  alkoxy,  $C_{1-6}$  haloalkoxy,  $SO_2R^3$ ,  $C_{1-6}$  hydroxyalkyl.

- 35  $R^3$  represents H or  $C_{1-6}$  alkyl;

or pharmaceutically acceptable salts, solvates or physiologically functional derivatives thereof in the preparation of a medicament for use in the treatment of a disorder mediated by inappropriate IKK3 activity.

5 In a sixth aspect there is provided a method of treating inflammatory and tissue repair disorders, particularly rheumatoid arthritis, inflammatory bowel disease, asthma and COPD (chronic obstructive pulmonary disease); osteoarthritis, osteoporosis and fibrotic diseases; dermatosis, including psoriasis, atopic dermatitis and ultraviolet radiation (UV)-  
10 induced skin damage; autoimmune diseases including systemic lupus erythematosus, multiple sclerosis, psoriatic arthritis, ankylosing spondylitis, tissue and organ rejection, Alzheimer's disease, stroke, atherosclerosis, restenosis, diabetes, glomerulonephritis, cancer, including Hodgkins disease, cachexia, inflammation associated with infection and certain viral infections, including acquired immune deficiency syndrome (AIDS), adult  
15 respiratory distress syndrome, and Ataxia Telangiectasia comprising administering a therapeutically effective amount to a mammal of a compound of Formula (Ia), or a salt, solvate or pharmaceutically functional derivative thereof.

In a seventh aspect there is provided the use of a compound of Formula (Ia) or a salt, solvate or physiologically functional derivative thereof in the manufacture of a medicament  
20 for the treatment of inflammatory and tissue repair disorders, particularly rheumatoid arthritis, inflammatory bowel disease, asthma and COPD (chronic obstructive pulmonary disease); osteoarthritis, osteoporosis and fibrotic diseases; dermatosis, including psoriasis, atopic dermatitis and ultraviolet radiation (UV)-induced skin damage; autoimmune diseases including systemic lupus erythematosus, multiple sclerosis, psoriatic  
25 arthritis, ankylosing spondylitis, tissue and organ rejection, Alzheimer's disease, stroke, atherosclerosis, restenosis, diabetes, glomerulonephritis, cancer, including Hodgkins disease, cachexia, inflammation associated with infection and certain viral infections, including acquired immune deficiency syndrome (AIDS), adult respiratory distress syndrome, and Ataxia Telangiectasia.

#### 30 DETAILED DESCRIPTION OF THE INVENTION

As used herein, the term "effective amount" means that amount of a drug or pharmaceutical agent that will elicit the biological or medical response of a tissue, system, animal or human that is being sought, for instance, by a researcher or clinician.

35 Furthermore, the term "therapeutically effective amount" means any amount which, as compared to a corresponding subject who has not received such amount, results in improved treatment, healing, prevention, or amelioration of a disease, disorder, or side effect, or a decrease in the rate of advancement of a disease or disorder. The term also includes within its scope amounts effective to enhance normal physiological function.

As used herein "a compound of the invention" or "a compound of Formula (I) or (Ia) means a compound of Formula (I) or (Ia) or a pharmaceutically acceptable salt, solvate or physiologically functional derivative thereof.

- 5 As used herein the term "alkyl" refers to a straight- or branched-chain hydrocarbon radical having the specified number of carbon atoms, so for example, as used herein, the terms "C<sub>1</sub>-C<sub>3</sub> alkyl" and "C<sub>1</sub>-C<sub>6</sub> alkyl" refer to an alkyl group, as defined above, containing at least 1, and at most 3 or 6 carbon atoms respectively. Examples of such branched or straight-chained alkyl groups useful in the present invention include, but are not limited to, methyl, ethyl, n-propyl, isopropyl, isobutyl, n-butyl, t-butyl, n-pentyl, isopentyl, and n-hexyl.

- 10 As used herein, the term "alkylene" refers to a straight or branched chain divalent hydrocarbon radical having the specified number of carbon atoms, so for example, as used herein, the terms "C<sub>1</sub>-C<sub>3</sub> alkylene" and "C<sub>1</sub>-C<sub>6</sub> alkylene" refer to an alkylene group, as defined above, which contains at least 1, and at most 3 or 6, carbon atoms respectively. Examples of "C<sub>1</sub>-C<sub>3</sub> alkylene" and "C<sub>1</sub>-C<sub>6</sub> alkylene" groups useful in the present invention include, but are not limited to, methylene, ethylene, n-propylene, n-butylene, isopentylene, and the like.

- 20 As used herein, the term "alkenyl" (and "alkenylene") refers to straight or branched hydrocarbon chains containing the specified number of carbon atoms and at least one and up to 3 carbon-carbon double bonds. Examples include ethenyl (and ethenylene) and propenyl (and propenylene).

- 25 As used herein, the term "halogen" refers to fluorine (F), chlorine (Cl), bromine (Br), or iodine (I) and the term "halo" refers to the halogen radicals: fluoro (-F), chloro (-Cl), bromo(-Br), and iodo(-I).

- 30 As used herein, the term "C<sub>1</sub>-C<sub>6</sub> haloalkyl" refers to a straight or branched chain alkyl group as defined above containing at least 1, and at most 6 carbon atoms respectively substituted with at least one halo group, halo being as defined herein. Examples of such branched or straight chained haloalkyl groups useful in the present invention include, but are not limited to, methyl, ethyl, propyl, isopropyl, isobutyl and n-butyl substituted independently with one or more halos, e.g., fluoro, chloro, bromo and iodo.

- 35 As used herein, the term "cycloalkyl" refers to a non-aromatic cyclic hydrocarbon ring containing the specified number of carbon atoms so, for example, the term "C<sub>5</sub>-C<sub>7</sub> cycloalkyl" refers to a non-aromatic cyclic hydrocarbon ring having from five to seven carbon atoms. Exemplary "C<sub>5</sub>-C<sub>7</sub> cycloalkyl" groups useful in the present invention include, but are not limited to, cyclopentyl, cyclohexyl and cycloheptyl.

As used herein, the term "cycloalkenyl" refers to a non-aromatic monocyclic carboxycyclic ring having the specified number of carbon atoms and up to 3 carbon-carbon double bonds. "Cycloalkenyl" includes by way of example cyclopentenyl and cyclohexenyl.

- 5 As used herein, the term "heterocyclic" or the term "heterocyclyl" refers to a non-aromatic heterocyclic ring containing the specified number ring atoms being saturated or having one or more degrees of unsaturation, containing one or more heteroatom substitutions selected from O and/or N. Such a ring may be optionally fused to one or more other "heterocyclic" ring(s) or cycloalkyl ring(s). Examples of "heterocyclic" moieties include, but are not limited to, tetrahydrofuran, pyran, 1,4-dioxane, 1,3-dioxane, piperidine, 10 piperazine, 2,4-piperazinedione, pyrrolidine, imidazolidine, pyrazolidine, morpholine, thiomorpholine, tetrahydrothiopyran, tetrahydrothiophene, and the like.

- 15 As used herein, the term "aryl" refers to monocyclic carbocyclic groups and fused bicyclic carbocyclic groups having the specified number of carbon atoms and having at least one aromatic ring. Examples of aryl groups include phenyl and naphthyl.

- 20 As used herein, the term "heteroaryl" refers to an aromatic monocyclic ring, or to a fused bicyclic or tricyclic ring system wherein at least one ring is aromatic, having the specified number of ring atoms and containing at least one heteroatom selected from N, O, and/or S. Examples of "heteroaryl" groups used herein include furanyl, thiophenyl, pyrrolyl, imidazolyl, pyrazolyl, triazolyl, tetrazolyl, thiazolyl, oxazolyl, isoxazolyl, oxadiazolyl, oxo-pyridyl, thiadiazolyl, isothiazolyl, pyridyl, pyridazyl, pyrazinyl, pyrimidyl, quinolinyl, 25 isoquinolinyl, benzofuranyl, benzothiophenyl, indolyl, indazolyl.

- 30 As used herein, the term "alkoxy" refers to the group  $R_aO-$ , where  $R_a$  is alkyl as defined above and the terms " $C_1-C_3$  alkoxy" and " $C_1-C_6$  alkoxy" refer to an alkoxy group as defined herein wherein the alkyl moiety contains at least 1, and at most 3 or 6, carbon atoms. Exemplary " $C_1-C_3$  alkoxy" and " $C_1-C_6$  alkoxy" groups useful in the present invention include, but are not limited to, methoxy, ethoxy, n-propoxy, isopropoxy, n-butoxy, and t-butoxy.

As used herein the term "hydroxyalkyl" refers to the group  $-R_aOH$ , where  $R_a$  is an alkylene group as defined above.

- 35 As used herein, the term "haloalkoxy" refers to the group  $R_aO-$ , where  $R_a$  is haloalkyl as defined above and the term " $C_1-C_6$  haloalkoxy" refers to a haloalkoxy group as defined herein wherein the haloalkyl moiety contains at least 1, and at most 6, carbon atoms. Exemplary  $C_1-C_6$  haloalkoxy groups useful in the present invention include, but is not limited to, trifluoromethoxy.

- 40 Preferably, n represents 1 or 2. More preferably, n represents 2. Preferably each  $R^1$  is in the 5, 6- position on the ring. When n is 2, preferably one  $R^1$  represents  $-OCH_3$ . More

preferably both R<sup>1</sup> represents -OCH<sub>3</sub>. Most preferably each R<sup>1</sup> is OCH<sub>3</sub> in the 5, 6 - position.

Preferably c is 1.

Preferably X<sup>1</sup> represents C<sub>1-6</sub> alkenyl.

Preferably Z<sup>1</sup> represents aryl or heteroaryl optionally substituted by one or more substituents independently selected from halogen, CF<sub>3</sub>, CH<sub>2</sub>OH, SO<sub>2</sub>CH<sub>3</sub>, CH<sub>3</sub>, -OCH<sub>3</sub>.

While the preferred groups for each variable have generally been listed above separately for each variable, preferred compounds of this invention includes those in which several or each variable in Formula (I) and (Ia) is selected from the preferred, more preferred, or most preferred groups for each variable. Therefore, this invention is intended to include all combinations of preferred, more preferred, and most preferred groups.

Specific examples of compounds of Formula (I) and (Ia) of the present invention include the following:

5-[5,6-bis(methyloxy)-1H-benzimidazol-1-yl]-3-({[4-(hydroxymethyl)phenyl]methyl}oxy)-2-thiophenecarbonitrile
5-[5,6-bis(methyloxy)-1H-benzimidazol-1-yl]-3-[(1R)-1-(2-chlorophenyl)ethyl]oxy}-2-thiophenecarbonitrile
5-(1H-benzimidazol-1-yl)-3-[(2-methylphenyl)methyl]oxy}-2-thiophenecarbonitrile
5-(1H-benzimidazol-1-yl)-3-[(phenylmethyl)oxy]-2-thiophenecarbonitrile
5-[5,6-bis(methyloxy)-1H-benzimidazol-1-yl]-3-[(1-methyl-1H-1,2,3-benzotriazol-6-yl)methyl]oxy}-2-thiophenecarbonitrile
5-(6-(methyloxy)-5-[[2-(4-morpholinyl)ethyl]oxy]-1H-benzimidazol-1-yl)-3-({[2-(trifluoromethyl)phenyl]methyl}oxy)-2-thiophenecarbonitrile
5-[5,6-bis(methyloxy)-1H-benzimidazol-1-yl]-3-[(2,5-difluorophenyl)methyl]oxy}-2-thiophenecarbonitrile
5-[5,6-bis(methyloxy)-1H-benzimidazol-1-yl]-3-({[2-(trifluoromethyl)phenyl]methyl}oxy)-2-thiophenecarbonitrile
5-[5,6-bis(methyloxy)-1H-benzimidazol-1-yl]-3-({[2-(methylsulfonyl)phenyl]methyl}oxy)-2-thiophenecarbonitrile
5-(5-chloro-1H-benzimidazol-1-yl)-3-[(phenylmethyl)oxy]-2-thiophenecarbonitrile
5-[5,6-bis(methyloxy)-1H-benzimidazol-1-yl]-3-[(2,6-difluorophenyl)methyl]oxy}-2-thiophenecarbonitrile
5-[5-[(3-hydroxypropyl)oxy]-6-(methyloxy)-1H-benzimidazol-1-yl]-3-({[2-(trifluoromethyl)phenyl]methyl}oxy)-2-thiophene-carbonitrile
5-[5,6-bis(methyloxy)-1H-benzimidazol-1-yl]-3-[(3-bromo-4-pyridinyl)methyl]oxy}-2-thiophenecarbonitrile

5-[5,6-bis(methyloxy)-1 <i>H</i> -benzimidazol-1-yl]-3-[(cyclohexylmethyl)oxy]-2-thiophenecarbonitrile
1-[5-cyano-4-({[2-(trifluoromethyl)phenyl]methyl}oxy)-2-thienyl]- <i>N</i> -[2-(4-morpholinyl)ethyl]-1 <i>H</i> -benzimidazole-5-carboxamide
5-[5,6-bis(methyloxy)-1 <i>H</i> -benzimidazol-1-yl]-3-({(1 <i>R</i> )-1-[2-(trifluoromethyl)phenyl]ethyl}oxy)-2-thiophenecarbonitrile
5-[5,6-bis(methyloxy)-1 <i>H</i> -benzimidazol-1-yl]-3-[(2-phenylethyl)oxy]-2-thiophenecarbonitrile
5-[5,6-bis(methyloxy)-1 <i>H</i> -benzimidazol-1-yl]-3-[(2-[(trifluoromethyl)oxy]phenyl)methyl]oxy]-2-thiophenecarbonitrile
5-[5,6-bis(methyloxy)-1 <i>H</i> -benzimidazol-1-yl]-3-({[2-(methyloxy)phenyl]methyl}oxy)-2-thiophenecarbonitrile
5-[5,6-bis(methyloxy)-1 <i>H</i> -benzimidazol-1-yl]-3-[(2-(4-morpholinyl)ethyl]oxy)-2-thiophenecarbonitrile
5-[5,6-bis(methyloxy)-1 <i>H</i> -benzimidazol-1-yl]-3-[(2-(2-oxo-1-pyrrolidinyl)ethyl]oxy)-2-thiophenecarbonitrile
5-[5,6-bis(methyloxy)-1 <i>H</i> -benzimidazol-1-yl]-3-[(tetrahydro-2-furanylmethyl)oxy]-2-thiophenecarbonitrile
5-[5,6-bis(methyloxy)-1 <i>H</i> -benzimidazol-1-yl]-3-[(tetrahydro-2 <i>H</i> -pyran-2-ylmethyl)oxy]-2-thiophenecarbonitrile
5-[5,6-bis(methyloxy)-1 <i>H</i> -benzimidazol-1-yl]-3-[(2-(phenyloxy)ethyl]oxy)-2-thiophenecarbonitrile
5-[5,6-bis(methyloxy)-1 <i>H</i> -benzimidazol-1-yl]-3-[(1 <i>S</i> )-1-(2-chlorophenyl)butyl]oxy)-2-thiophenecarbonitrile
5-[5,6-bis(methyloxy)-1 <i>H</i> -benzimidazol-1-yl]-3-[(3-thienylmethyl)oxy]-2-thiophenecarbonitrile
5-[5,6-bis(methyloxy)-1 <i>H</i> -benzimidazol-1-yl]-3-[(2-thienylmethyl)oxy]-2-thiophenecarbonitrile
5-[5,6-bis(methyloxy)-1 <i>H</i> -benzimidazol-1-yl]-3-[(1 <i>R</i> )-1-methylpropyl]oxy)-2-thiophenecarbonitrile

As used herein, the term "optionally" means that the subsequently described event(s) may or may not occur, and includes both event(s), which occur, and events that do not occur.

- 5 As used herein, the term "physiologically functional derivative" refers to any pharmaceutically acceptable derivative of a compound of the present invention, for example, an ester or an amide, which upon administration to a mammal is capable of providing (directly or indirectly) a compound of the present invention or an active metabolite thereof. Such derivatives are clear to those skilled in the art, without undue experimentation, and with reference to the teaching of Burger's Medicinal Chemistry And
- 10 Drug Discovery, 5<sup>th</sup> Edition, Vol 1: Principles and Practice, which is incorporated herein by reference to the extent that it teaches physiologically functional derivatives.

As used herein, the term "solvate" refers to a complex of variable stoichiometry formed by a solute (in this invention, a compound of Formula (I) or (Ia) or a salt or physiologically functional derivative thereof) and a solvent. Such solvents for the purpose of the invention may not interfere with the biological activity of the solute. Examples of suitable solvents include, but are not limited to, water, methanol, ethanol and acetic acid. Preferably the solvent used is a pharmaceutically acceptable solvent. Examples of suitable pharmaceutically acceptable solvents include, without limitation, water, ethanol and acetic acid. Most preferably the solvent used is water.

As used herein, the term "substituted" refers to substitution with the named substituent or substituents, multiple degrees of substitution being allowed unless otherwise stated.

Certain of the compounds described herein may contain one or more chiral atoms, or may otherwise be capable of existing as two enantiomers. The compounds of this invention include mixtures of enantiomers as well as purified enantiomers or enantiomerically enriched mixtures. Also included within the scope of the invention are the individual isomers of the compounds represented by Formula (I) or (Ia) above as well as any wholly or partially equilibrated mixtures thereof. The present invention also covers the individual isomers of the compounds represented by the formulas above as mixtures with isomers thereof in which one or more chiral centers are inverted. Also, it is understood that any tautomers and mixtures of tautomers of the compounds of Formula (I) or (Ia) are included within the scope of the compounds of Formula (I) or (Ia).

The present invention also covers salts of the compounds of Formula (I) or (Ia). Typically, the salts of the present invention are pharmaceutically acceptable salts. Salts encompassed within the term "pharmaceutically acceptable salts" refer to non-toxic salts of the compounds of this invention. Salts of the compounds of the present invention may comprise acid addition salts derived from a nitrogen on a substituent in the compound of Formula (I). Representative salts include the following salts: acetate, benzenesulfonate, benzoate, bicarbonate, bisulfate, bitartrate, borate, bromide, calcium edetate, camsylate, carbonate, chloride, clavulanate, citrate, dihydrochloride, edetate, edisylate, estolate, esylate, fumarate, gluceptate, gluconate, glutamate, glycolylarsanilate, hexylresorcinate, hydrabamine, hydrobromide, hydrochloride, hydroxynaphthoate, iodide, isethionate, lactate, lactobionate, laurate, malate, maleate, mandelate, mesylate, methylbromide, methylnitrate, methylsulfate, monopotassium maleate, mucate, napsylate, nitrate, N-methylglucamine, oxalate, palmoate (embonate), palmitate, pantothenate, phosphate/diphosphate, polygalacturonate, potassium, salicylate, sodium, stearate, subacetate, succinate, tannate, tartrate, teoclate, tosylate, triethiodide, trimethylammonium and valerate. Other salts, which are not pharmaceutically acceptable, may be useful in the preparation of compounds of this invention and these form a further aspect of the invention.



While it is possible that, for use in therapy, therapeutically effective amounts of a compound of Formula (I) or (Ia), as well as salts, solvates and physiological functional derivatives thereof, may be administered as the raw chemical, it is possible to present the active ingredient as a pharmaceutical composition. Accordingly, the invention further provides pharmaceutical compositions, which include therapeutically effective amounts of compounds of the Formula (I) or (Ia) and salts, solvates and physiological functional derivatives thereof, and one or more pharmaceutically acceptable carriers, diluents, or excipients. The compounds of the Formula (I) or (Ia) and salts, solvates and physiological functional derivatives thereof, are as described above. The carrier(s), diluent(s) or excipient(s) must be acceptable in the sense of being compatible with the other ingredients of the formulation and not deleterious to the recipient thereof. In accordance with another aspect of the invention there is also provided a process for the preparation of a pharmaceutical formulation including admixing a compound of the Formula (I) or (Ia), or salts, solvates and physiological functional derivatives thereof, with one or more pharmaceutically acceptable carriers, diluents or excipients.

Pharmaceutical formulations may be presented in unit dose forms containing a predetermined amount of active ingredient per unit dose. Such a unit may contain, for example, 0.5mg to 1g, preferably 1mg to 700mg, more preferably 5mg to 100mg of a compound of the Formula (I), depending on the condition being treated, the route of administration and the age, weight and condition of the patient, or pharmaceutical formulations may be presented in unit dose forms containing a predetermined amount of active ingredient per unit dose. Preferred unit dosage formulations are those containing a daily dose or sub-dose, as herein above recited, or an appropriate fraction thereof, of an active ingredient. Furthermore, such pharmaceutical formulations may be prepared by any of the methods well known in the pharmacy art.

Pharmaceutical formulations may be adapted for administration by any appropriate route, for example by the oral (including buccal or sublingual), rectal, nasal, topical (including buccal, sublingual or transdermal), vaginal or parenteral (including subcutaneous, intramuscular, intravenous or intradermal) route. Such formulations may be prepared by any method known in the art of pharmacy, for example by bringing into association the active ingredient with the carrier(s) or excipient(s).

Pharmaceutical formulations adapted for oral administration may be presented as discrete units such as capsules or tablets; powders or granules; solutions or suspensions in aqueous or non-aqueous liquids; edible foams or whips; or oil-in-water liquid emulsions or water-in-oil liquid emulsions.

For instance, for oral administration in the form of a tablet or capsule, the active drug component can be combined with an oral, non-toxic pharmaceutically acceptable inert

carrier such as ethanol, glycerol, water and the like. Powders are prepared by comminuting the compound to a suitable fine size and mixing with a similarly comminuted pharmaceutical carrier such as an edible carbohydrate, as, for example, starch or mannitol. Flavoring, preservative, dispersing and coloring agent can also be present.

5

Capsules are made by preparing a powder mixture, as described above, and filling formed gelatin sheaths. Glidants and lubricants such as colloidal silica, talc, magnesium stearate, calcium stearate or solid polyethylene glycol can be added to the powder mixture before the filling operation. A disintegrating or solubilizing agent such as agar-agar, calcium carbonate or sodium carbonate can also be added to improve the availability of the medicament when the capsule is ingested.

10

Moreover, when desired or necessary, suitable binders, lubricants, disintegrating agents and coloring agents can also be incorporated into the mixture. Suitable binders include starch, gelatin, natural sugars such as glucose or beta-lactose, corn sweeteners, natural and synthetic gums such as acacia, tragacanth or sodium alginate, carboxymethylcellulose, polyethylene glycol, waxes and the like. Lubricants used in these dosage forms include sodium oleate, sodium stearate, magnesium stearate, sodium benzoate, sodium acetate, sodium chloride and the like. Disintegrators include, without limitation, starch, methyl cellulose, agar, bentonite, xanthan gum and the like. Tablets are formulated, for example, by preparing a powder mixture, granulating or slugging, adding a lubricant and disintegrant and pressing into tablets. A powder mixture is prepared by mixing the compound, suitably comminuted, with a diluent or base as described above, and optionally, with a binder such as carboxymethylcellulose, an aliginate, gelatin, or polyvinyl pyrrolidone, a solution retardant such as paraffin, a resorption accelerator such as a quaternary salt and/or an absorption agent such as bentonite, kaolin or dicalcium phosphate. The powder mixture can be granulated by wetting with a binder such as syrup, starch paste, acadia mucilage or solutions of cellulosic or polymeric materials and forcing through a screen. As an alternative to granulating, the powder mixture can be run through the tablet machine and the result is imperfectly formed slugs broken into granules. The granules can be lubricated to prevent sticking to the tablet forming dies by means of the addition of stearic acid, a stearate salt, talc or mineral oil. The lubricated mixture is then compressed into tablets. The compounds of the present invention can also be combined with a free flowing inert carrier and compressed into tablets directly without going through the granulating or slugging steps. A clear or opaque protective coating consisting of a sealing coat of shellac, a coating of sugar or polymeric material and a polish coating of wax can be provided. Dyestuffs can be added to these coatings to distinguish different unit dosages.

25

30

35

Oral fluids such as solution, syrups and elixirs can be prepared in dosage unit form so that a given quantity contains a predetermined amount of the compound. Syrups can be prepared by dissolving the compound in a suitably flavored aqueous solution, while elixirs

40

are prepared through the use of a non-toxic alcoholic vehicle. Suspensions can be formulated by dispersing the compound in a non-toxic vehicle. Solubilizers and emulsifiers such as ethoxylated isostearyl alcohols and polyoxy ethylene sorbitol ethers, preservatives, flavor additive such as peppermint oil or natural sweeteners or saccharin or other artificial sweeteners, and the like can also be added.

Where appropriate, dosage unit formulations for oral administration can be microencapsulated. The formulation can also be prepared to prolong or sustain the release as for example by coating or embedding particulate material in polymers, wax or the like.

The compounds of Formula (I) or (Ia), and salts, solvates and physiological functional derivatives thereof, can also be administered in the form of liposome delivery systems, such as small unilamellar vesicles, large unilamellar vesicles and multilamellar vesicles. Liposomes can be formed from a variety of phospholipids, such as cholesterol, stearylamine or phosphatidylcholines.

The compounds of Formula (I) or (Ia) and salts, solvates and physiological functional derivatives thereof may also be delivered by the use of monoclonal antibodies as individual carriers to which the compound molecules are coupled. The compounds may also be coupled with soluble polymers as targetable drug carriers. Such polymers can include polyvinylpyrrolidone, pyran copolymer, polyhydroxypropylmethacrylamide -phenol, polyhydroxyethylaspartamidephenol, or polyethyleneoxidepolylysine substituted with palmitoyl residues. Furthermore, the compounds may be coupled to a class of biodegradable polymers useful in achieving controlled release of a drug, for example, polylactic acid, polepsilon caprolactone, polyhydroxy butyric acid, polyorthoesters, polyacetals, polydihydropyrans, polycyanoacrylates and cross-linked or amphipathic block copolymers of hydrogels.

Pharmaceutical formulations adapted for transdermal administration may be presented as discrete patches intended to remain in intimate contact with the epidermis of the recipient for a prolonged period of time. For example, the active ingredient may be delivered from the patch by iontophoresis as generally described in Pharmaceutical Research, 3(6), 318 (1986).

Pharmaceutical formulations adapted for topical administration may be formulated as ointments, creams, suspensions, lotions, powders, solutions, pastes, gels, sprays, aerosols or oils.

For treatments of the eye or other external tissues, for example mouth and skin, the formulations are preferably applied as a topical ointment or cream. When formulated in an ointment, the active ingredient may be employed with either a paraffinic or a water-

miscible ointment base. Alternatively, the active ingredient may be formulated in a cream with an oil-in-water cream base or a water-in-oil base.

5      Pharmaceutical formulations adapted for topical administrations to the eye include eye drops wherein the active ingredient is dissolved or suspended in a suitable carrier, especially an aqueous solvent.

10      Pharmaceutical formulations adapted for topical administration in the mouth include lozenges, pastilles and mouth washes.

10      Pharmaceutical formulations adapted for rectal administration may be presented as suppositories or as enemas.

15      Pharmaceutical formulations adapted for nasal administration wherein the carrier is a solid include a coarse powder having a particle size for example in the range 20 to 500 microns which is administered in the manner in which snuff is taken, i.e. by rapid inhalation through the nasal passage from a container of the powder held close up to the nose. Suitable formulations wherein the carrier is a liquid, for administration as a nasal spray or as nasal drops, include aqueous or oil solutions of the active ingredient.

20      Pharmaceutical formulations adapted for administration by inhalation include fine particle dusts or mists, which may be generated by means of various types of metered, dose pressurised aerosols, nebulizers or insufflators.

25      Pharmaceutical formulations adapted for vaginal administration may be presented as pessaries, tampons, creams, gels, pastes, foams or spray formulations.

30      Pharmaceutical formulations adapted for parenteral administration include aqueous and non-aqueous sterile injection solutions which may contain anti-oxidants, buffers, bacteriostats and solutes which render the formulation isotonic with the blood of the intended recipient; and aqueous and non-aqueous sterile suspensions which may include suspending agents and thickening agents. The formulations may be presented in unit-dose or multi-dose containers, for example sealed ampoules and vials, and may be stored in a freeze-dried (lyophilized) condition requiring only the addition of the sterile liquid carrier, for example water for injections, immediately prior to use. Extemporaneous  
35      injection solutions and suspensions may be prepared from sterile powders, granules and tablets.

40      It should be understood that in addition to the ingredients particularly mentioned above, the formulations may include other agents conventional in the art having regard to the type of formulation in question, for example those suitable for oral administration may include flavouring agents.

A therapeutically effective amount of a compound of the present invention will depend upon a number of factors including, for example, the age and weight of the animal, the precise condition requiring treatment and its severity, the nature of the formulation, and the route of administration, and will ultimately be at the discretion of the attendant physician or veterinarian. However, an effective amount of a compound of Formula (I) or (Ia) will generally be in the range of 0.1 to 100 mg/kg body weight of recipient (mammal) per day and more usually in the range of 1 to 10 mg/kg body weight per day. Thus, for a 70kg adult mammal, the actual amount per day would usually be from 70 to 700 mg and this amount may be given in a single dose per day or more usually in a number (such as two, three, four, five or six) of sub-doses per day such that the total daily dose is the same. An effective amount of a salt or solvate, or physiologically functional derivative thereof, may be determined as a proportion of the effective amount of the compound of Formula (I) or (Ia) *per se*.

The compounds of Formula (I) or (Ia) and salts, solvates and physiological functional derivatives thereof, are believed to have utility in as a result of inhibition of the protein kinase IKK3, in particular in the treatment and prevention of disease states mediated by IKK3 mechanisms including inflammatory and tissue repair disorders, particularly rheumatoid arthritis, inflammatory bowel disease, asthma and COPD (chronic obstructive pulmonary disease); osteoarthritis, osteoporosis and fibrotic diseases; dermatosis, including psoriasis, atopic dermatitis and ultraviolet radiation (UV)-induced skin damage; autoimmune diseases including systemic lupus erythematosus, multiple sclerosis, psoriatic arthritis, ankylosing spondylitis, tissue and organ rejection, Alzheimer's disease, stroke, atherosclerosis, restenosis, diabetes, glomerulonephritis, cancer, including Hodgkins disease, cachexia, inflammation associated with infection and certain viral infections, including acquired immune deficiency syndrome (AIDS), adult respiratory distress syndrome, and Ataxia Telangiectasia.

The present invention thus also provides compounds of Formula (I) or (Ia) and pharmaceutically acceptable salts or solvates thereof, or physiologically functional derivatives thereof, for use in medical therapy, and particularly in the treatment of disorders mediated by IKK3 activity.

The inappropriate IKK3 activity referred to herein is any IKK3 activity that deviates from the normal IKK3 activity expected in a particular mammalian subject. Inappropriate IKK3 activity may take the form of, for instance, an abnormal increase in activity, or an aberration in the timing and or control of IKK3 activity. Such inappropriate activity may result then, for example, from overexpression or mutation of the protein kinase leading to inappropriate or uncontrolled activation.

The present invention is directed to methods of regulating, modulating, or inhibiting IKK3 for the prevention and/or treatment of disorders related to unregulated IKK3 activity. In particular, the compounds of the present invention can also be used in inflammatory disease, particularly rheumatoid arthritis.

5 A further aspect of the invention provides a method of treatment of a mammal suffering from a disorder mediated by IKK3 activity, which includes administering to said subject an effective amount of a compound of Formula (I) or a pharmaceutically acceptable salt, solvate, or a physiologically functional derivative thereof.

10 A further aspect of the present invention provides the use of a compound of Formula (I), or a pharmaceutically acceptable salt or solvate thereof, or a physiologically functional derivative thereof, in the preparation of a medicament for the treatment of a disorder characterized by inappropriate IKK3 activity.


15 Particular disorders characterised by inappropriate IKK3 activity in particular in the treatment and prevention of disease states mediated by IKK3 mechanisms including inflammatory and tissue repair disorders, particularly rheumatoid arthritis, inflammatory bowel disease, asthma and COPD (chronic obstructive pulmonary disease); osteoarthritis, osteoporosis and fibrotic diseases; dermatosis, including psoriasis, atopic dermatitis and  
20 ultraviolet radiation (UV)-induced skin damage; autoimmune diseases including systemic lupus erythematosus, multiple sclerosis, psoriatic arthritis, ankylosing spondylitis, tissue and organ rejection, Alzheimer's disease, stroke, atherosclerosis, restenosis, diabetes, glomerulonephritis, cancer, including Hodgkins disease, cachexia, inflammation  
25 associated with infection and certain viral infections, including acquired immune deficiency syndrome (AIDS), adult respiratory distress syndrome, and Ataxia Telangiectasia.  
as a result of inhibition of the protein kinase IKK3.

30 The compounds of this invention may be made by a variety of methods, including standard chemistry. Any previously defined variable will continue to have the previously defined meaning unless otherwise indicated. Illustrative general synthetic methods are set out below and then specific compounds of the invention are prepared in the Working Examples.


35 Compounds of general Formula (I) and (Ia) may be prepared by methods known in the art of organic synthesis as set forth in part by the following synthesis schemes. In all of the schemes described below, it is well understood that protecting groups for sensitive or reactive groups are employed where necessary in accordance with general principles of chemistry. Protecting groups are manipulated according to standard methods of organic  
40 synthesis (T. W. Green and P. G. M. Wuts (1991) Protecting Groups in Organic Synthesis, John Wiley & Sons). These groups are removed at a convenient stage of the compound synthesis using methods that are readily apparent to those skilled in the art. The selection

of processes as well as the reaction conditions and order of their execution shall be consistent with the preparation of compounds of Formula (I) and (Ia). Those skilled in the art will recognize if a stereocenter exists in compounds of Formula (I) and (Ia). Accordingly, the present invention includes both possible stereoisomers and includes not only racemic compounds but the individual enantiomers as well. When a compound is desired as a single enantiomer, it may be obtained by stereospecific synthesis or by resolution of the final product or any convenient intermediate. Resolution of the final product, an intermediate, or a starting material may be effected by any suitable method known in the art. See, for example, Stereochemistry of Organic Compounds by E. L. Eliel, S. H. Wilen, and L. N. Mander (Wiley-Interscience, 1994).

Compounds of Formula I and (Ia) can be prepared for example according to the general scheme depicted below and the Examples section following:




A compound of Formula (1) or (Ia) may be prepared by dehydration of a Compound of Formula (II)



wherein all variables are as defined for a compound of Formula (I) and (Ia).

The dehydration reaction can be carried out using a variety of reagents. Suitable dehydration reagents include but are not limited to thionyl chloride, trifluoroacetic anhydride, phosphorous oxychloride, phosphorous pentoxide, and *N,N*-dicyclohexylcarbodiimide and 2-chloro-1,3-dimethylimidazolinium chloride. The reaction may be optionally heated to from about 50 to about 150 °C. Suitable solvents for this reaction include but are not limited to dichloromethane, chloroform, benzene, toluene, *N,N*-dimethylformamide and dichloroethane.


A compound of Formula (II) may be prepared directly from a compound of Formula (III).



wherein all variables are as defined in connection with Formula (1) and (Ia).  $R^4$  represents for example, a suitable carboxylic acid protecting group,  $C_{1-6}$  alkyl,  $C_{1-6}$  alkenyl,  $C_{1-6}$  alkynyl or  $C_{5-7}$  cycloalkyl.

This reaction is typically performed in a sealed vessel with an excess of ammonia. The reaction is typically heated to a temperature of from about 50 to about 120 °C. Suitable solvents for this reaction include but are not limited to methanol, ethanol, isopropanol, tetrahydrofuran, and dioxane.

A compound of Formula (III) may be prepared from compounds of Formula (IV):



LG is a suitable leaving group; and all other variables are as defined in connection with Scheme 1 above.

In general the process for preparing a compound of Formula (III) comprises the steps of:




- a) reacting the compound of Formula (IV) with a base and a compound of Formula (XII) to prepare a compound of the Formula (III); or  
 b) reacting the compound of Formula (IV) with a compound of Formula (XIII) under Mitsunobu conditions to prepare a compound of Formula (III).

More specifically, a compound of Formula (III) can be prepared by reacting a compound of Formula (IV) with a compound of Formula (XII). The compounds of Formula (XII) are commercially available or can be prepared using conventional knowledge in the art. The reaction may be carried out in an inert solvent, conveniently at room temperature, in the presence of a suitable base. The compound of Formula (IV) and the compound of Formula (XII) may be present in equimolar amounts; however, a slight excess of the compound of Formula (XII) may be employed if desired. Examples of suitable bases for this reaction include but are not limited to, potassium carbonate, sodium carbonate, cesium carbonate, sodium hydride, and potassium hydride. Examples of suitable inert solvents for this reaction include but are not limited to, *N,N*-dimethylformamide, tetrahydrofuran, dioxane, and 1,2-dimethoxyethane.

In another embodiment, a compound of Formula (III) can be prepared by reacting a compound of Formula (IV) with a compound of Formula (XIII). The compounds of Formula (XIII) are commercially available or can be prepared using conventional knowledge in the art. The reaction is carried out in an inert solvent under standard Mitsunobu conditions. See, Hughes, D.L., *Org. React.* **42**:335-656 (1992); and Mitsunobu, O., *Synthesis* 1-28 (1981). Typically the compound of Formula (III), the compound of Formula (XII), a triarylphosphine, and a dialkyl azodicarboxylate are reacted together at room temperature. Examples of suitable triarylphosphines include but are not limited to, triphenylphosphine, tri-*p*-tolylphosphine, and trimesitylphosphine. Examples of suitable dialkyl azodicarboxylates include but are not limited to, diethyl azodicarboxylate, diisopropyl azodicarboxylate, and di-*tert*-butyl azodicarboxylate. Examples of suitable inert solvents for this reaction include but are not limited to, tetrahydrofuran, dioxane, 1,2-dimethoxyethane, dichloromethane, and toluene.


More specifically, compounds of Formula (IV) can be prepared by reacting a compound of Formula (VII) with a compound of Formula (V) to prepare a compound of Formula (IV).



wherein all variables are as defined as previously.

The reaction of a compound of Formula (VII) with a compound of Formula (V) is typically carried out in an inert solvent at room temperature. Typically two molar equivalents of a compound of Formula (VII) are combined with one molar equivalent of a compound of Formula (V). Examples of suitable inert solvents for this reaction include but are not limited to, chloroform, dichloromethane, tetrahydrofuran, dioxane, and toluene.

A compound of Formula (V) can be prepared by reacting a compound of Formula (VI) with sulfuryl chloride.




wherein all variables are as defined as previously.

Compounds of Formula (VI) are commercially available or can be prepared using conventional knowledge in the art. Typically, reaction of a compound of Formula (VI) with sulfuryl chloride at room temperature provides a compound of Formula (V). Excess sulfuryl chloride may be used if desired. Examples of suitable solvents include but are not limited to chloroform, dichloromethane, and toluene. See, Corral, C.; Lissavetzky, J. *Synthesis* 847-850 (1984).

A compound of Formula VII may be prepared according to the process outlined in the Scheme below:

## Scheme 2




wherein all variables are as defined as previously.

Generally, this process for preparing the a compound of Formula (VII) (all formulas and all variables having been defined above comprises the steps of:

- a) reacting a compound of Formula (XI) with a suitable acylating agent to prepare a compound of the Formula (X);
- b) reacting a compound of Formula (X) under nitration conditions to prepare a compound of the Formula (IX);
- c) reducing a compound of Formula (IX) to prepare a compound of Formula (VIII);
- and
- d) cyclizing a compound of Formula (VIII) to prepare a compound of Formula (VII).

The order of the foregoing steps is not critical to the practice of the invention and the process may be practiced by performing the steps in any suitable order based on the knowledge of those skilled in the art.

More specifically, a compound of Formula (VII) can be prepared by cyclizing a compound of Formula (VIII):




wherein all variables are as defined above.

This type of cyclization reaction is well documented in the literature. See, Braña, M. F., et. al., *J. Med. Chem.* **45**: 5813-5816 (2002); Fonseca, T., et. al., *Tetrahedron* **57**: 1793-1799

(2001); White, A. W., et. al., *J. Med. Chem.* **43**: 4084-4097 (2000); and Tamura, S. Y., et. al., *Bioorg. Med. Chem. Lett.* **7**: 1359-1364 (1997). This reaction may be carried out neat or in a suitable solvent. The reaction may optionally be heated to a temperature of from about 50 to about 200 °C. Typically an excess of a suitable acid is used. Examples of suitable acids include but are not limited to acetic acid, trifluoroacetic acid, hydrochloric acid, hydrobromic acid, sulfuric acid, methanesulfonic acid, p-toluenesulfonic acid, and pyridinium p-toluenesulfonate. A dehydrating reagent may optionally be used as well. Examples of suitable dehydrating reagents include but are not limited to magnesium sulfate, sodium sulfate, phosphorous pentoxide, and molecular sieves. Examples of suitable solvents include but are not limited to dichloromethane, chloroform, toluene, xylenes, methanol, ethanol, and water.


A compound of Formula (VIII) may be prepared by reducing a compound of Formula (IX).



wherein all variables are as defined above.

The reduction can be carried out using conventional techniques and reducing agents. See, Rangarajan, M., et al., *Bioorg. Med. Chem.* **8**:2591-2600 (2000); White, A.W., et al., *J. Med. Chem.* **43**: 4084-4097 (2000); Silvestri, R., et al., *Bioorg. Med. Chem.* **8**:2305-2309 (2000); Nagaraja, D., et al., *Tetrahedron Lett.* **40**:7855-7856 (1999); Jung, F., et al., *J. Med. Chem.* **34**:1110-1116 (1991); Srivastava, R.P., et al., *Pharmazie* **45**:34-37 (1990); Hankovszky, H.O., et al., *Can. J. Chem.* **67**:1392-1400 (1989); Ladd, D.L., et al., *J. Org. Chem.* **53**:417-420 (1988); Mertens, A., et al., *J. Med. Chem.* **30**:1279-1287 (1987); and Sharma, K.S., et al., *Synthesis* **4**:316-318 (1981). Examples of suitable reducing agents for this reaction include but are not limited to, palladium with hydrogen, palladium with ammonium formate, platinum oxide with hydrogen, nickel with hydrogen, tin(II) chloride, iron with acetic acid, aluminum with ammonium chloride, borane, sodium dithionite, and hydrazine. The reaction may optionally be heated to between about 50 and about 120 °C. Suitable solvents for this reaction vary and include but are not limited to, water, methanol, ethanol, ethyl acetate, tetrahydrofuran, and dioxane.

A compound of Formula (IX) may be prepared by reacting a compound of Formula (X) under nitration conditions.



wherein all variables are as defined above.

The compound of Formula (IX) can be prepared by reacting a compound of Formula (X) under nitration conditions.

The protection of anilines e.g. in Formula (X) by the moiety  $-\text{COR}^5$  (wherein  $\text{R}^5$  is a protecting group not H as depicted above) is a common transformation well known to one skilled in the art. See, Kocienski, P.J. *Protecting Groups*, Georg Thieme Verlag, Stuttgart, 1994; and Greene, T.W., Wuts, P. G. M. *Protecting Groups in Organic Synthesis* (2<sup>nd</sup> Edition), J. Wiley and Sons, 1991. Suitable protecting groups for this application include but are not limited to acetyl, trifluoroacetyl, benzyloxycarbonyl, allyloxycarbonyl, 2-(trimethylsilyl)ethoxycarbonyl, phenylsulfonyl, and p-toluenesulfonyl. Reagents and conditions vary according to the nature of the particular protecting group. Some typical reagents include but are not limited to acetic anhydride, trifluoroacetic anhydride, benzyl chloroformate, allyl chloroformate, 4-nitrophenyl 2-(trimethylsilyl)ethyl carbonate, phenylsulfonyl chloride, and p-toluenesulfonyl chloride. In certain cases the addition of some base is required. Examples of suitable bases include but are not limited to potassium carbonate, sodium carbonate, trialkylamines, pyridine, and potassium t-butoxide. Suitable solvents for these conversions include but are not limited to dichloromethane, chloroform, tetrahydrofuran, acetic acid, methanol, ethanol, water, toluene, and diethyl ether.


The nitration of anilines is also well documented in the literature and the foregoing reaction may be carried out using these conventional techniques. See, Wissner, A., et. al., *J. Med. Chem.* **46**: 49-63 (2003); Duggan, S. A., et. al., *J. Org. Chem.* **66**: 4419-4426 (2001); Clews, J., et. al., *Tetrahedron* **56**: 8735-8746 (2000); and Kagechika, H., *J. Med. Chem.* **31**: 2182-2192 (1988). The nitration may be carried out with a variety of nitrating reagents including but not limited to 70% aqueous nitric acid, red fuming nitric acid, ammonium nitrate with trifluoroacetic anhydride, and potassium nitrate with trifluoromethanesulfonic acid. The reaction is typically conducted at room temperature, but may be optionally heated to a temperature of from about 40 to about 100 °C in certain cases. Suitable solvents include but are not limited to acetic acid, sulfuric acid, acetic anhydride, dichloromethane, and chloroform.

The nitration results in a compound of Formula (IX), Formula (VII) can be accomplished through many different conventional methods. See, Kocienski, P.J. *Protecting Groups*, Georg Thieme Verlag, Stuttgart, 1994; and Greene, T.W., Wuts, P. G. M. *Protecting Groups in Organic Synthesis (2<sup>nd</sup> Edition)*, J. Wiley and Sons, 1991.

5

A compound of Formula (X) may be prepared by acylating a compound of Formula (XI).

wherein all variables are as defined above.



- 10 Acylation of anilines is a common transformation well known to one skilled in the art and such conventional acylation techniques may be employed for carrying out the foregoing reaction. See, Larock, R. C. *Comprehensive Organic Transformations*, VCH Publishers, Inc., New York, pp. 972-976, 979, 981 (1989). The acylation reaction is typically carried out using an acylating agent such as an acid halide, acid anhydride, or carboxylic acid, in
- 15 the presence of a coupling reagent(s). Examples of suitable coupling reagents include but are not limited to *N,N'*-dicyclohexylcarbodiimide, 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride, *O*-(7-azabenzotriazol-1-yl)-*N,N,N',N'*-tetramethyluronium hexafluorophosphate, and *N,N'*-carbonyldiimidazole. Suitable solvents include but are not limited to *N,N*-dimethylformamide, tetrahydrofuran, dioxane, toluene, benzene, 1,2-
- 20 dimethoxyethane, and 1-methyl-2-pyrrolidinone. Anilines of Formula (XI) are commercially available or readily prepared from commercially available material using conventional techniques.

- 25 Reaction conditions will be apparent to a skilled person and Examples of suitable conditions are further exemplified in the Examples Section below. It will also be evident from the above scheme that a compound of Formula (I) may be converted into another compound of Formula (I) using conventional procedures.

- 30 Certain embodiments of the present invention will now be illustrated by way of example only. The physical data given for the compounds exemplified is consistent with the assigned structure of those compounds.

## EXAMPLES

- 35 As used herein the symbols and conventions used in these processes, schemes and examples are consistent with those used in the contemporary scientific literature, for example, the *Journal of the American Chemical Society* or the *Journal of Biological*

*Chemistry.* Standard single-letter or three-letter abbreviations are generally used to designate amino acid residues, which are assumed to be in the L-configuration unless otherwise noted. Unless otherwise noted, all starting materials were obtained from commercial suppliers and used without further purification. Specifically, the following

5 abbreviations may be used in the examples and throughout the specification:

	g (grams);	mg (milligrams);
	L (liters);	mL (milliliters);
	μL (microliters);	psi (pounds per square inch);
10	M (molar);	mM (millimolar);
	i. v. (intravenous);	Hz (Hertz);
	MHz (megahertz);	mol (moles);
	mmol (millimoles);	rt (room temperature);
	min (minutes);	h (hours);
15	mp (melting point);	TLC (thin layer chromatography);
	T <sub>R</sub> (retention time);	RP (reverse phase);
	MeOH (methanol);	<i>i</i> -PrOH (isopropanol);
	TEA (triethylamine);	TFA (trifluoroacetic acid);
	TFAA (trifluoroacetic anhydride);	THF (tetrahydrofuran);
20	DMSO (dimethylsulfoxide);	AcOEt (ethyl acetate);
	DME (1,2-dimethoxyethane);	DCM (dichloromethane);
	DCE (dichloroethane);	DMF ( <i>N,N</i> -dimethylformamide);
	DMPU ( <i>N,N'</i> -dimethylpropyleneurea);	CDI (1,1-carbonyldiimidazole);
	IBCF (isobutyl chloroformate);	HOAc (acetic acid);
25	HOSu ( <i>N</i> -hydroxysuccinimide);	HOBT (1-hydroxybenzotriazole);
	mCPBA (meta-chloroperbenzoic acid);	
	EDC (1-[3-dimethylamino] propyl]-3-ethylcarbodiimide hydrochloride);	
	BOC ( <i>tert</i> -butoxycarbonyl);	Fmoc (9-fluorenylmethoxycarbonyl);
	DCC (dicyclohexylcarbodiimide);	CBZ (benzyloxycarbonyl);
30	Ac (acetyl);	atm (atmosphere);
	TMSE (2-(trimethylsilyl)ethyl);	TMS (trimethylsilyl);
	TIPS (triisopropylsilyl);	TBS ( <i>t</i> -butyldimethylsilyl);
	DMAP (4-dimethylaminopyridine);	BSA (bovine serum albumin)
	ATP (adenosine triphosphate);	HRP (horseradish peroxidase);
35	DMEM (Dulbecco's modified Eagle medium);	
	HPLC (high pressure liquid chromatography);	
	BOP (bis(2-oxo-3-oxazolidinyl)phosphinic chloride);	
	TBAF (tetra- <i>n</i> -butylammonium fluoride);	
	HBTU(O-Benzotriazole-1-yl- <i>N,N,N',N'</i> -tetramethyluroniumhexafluoro phosphate).	
40	HEPES (4-(2-hydroxyethyl)-1-piperazine ethane sulfonic acid);	
	DPPA (diphenylphosphoryl azide);	
	fHNO <sub>3</sub> (fuming HNO <sub>3</sub> ); and	

EDTA (ethylenediaminetetraacetic acid).  
 TMEDA (N,N,N',N'-tetramethyl-1,2-ethanediamine)  
 NBS (N-bromosuccinimide)  
 HATU (O-(7-azabenzobenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium  
 5 hexafluorophosphate)  
 DIPEA (diisopropylethylamine)  
 dppf (1,1'-bis(diphenylphosphino)ferrocene)  
 NIS (N-iodosuccinimide)  
 PTFE ((poly)tetrafluoroethylene)  
 10 LC-MS (liquid chromatography – mass spectrometry)

All references to ether are to diethyl ether; brine refers to a saturated aqueous solution of NaCl. Unless otherwise indicated, all temperatures are expressed in °C (degrees Centigrade). All reactions are conducted under an inert atmosphere at room temperature  
 15 unless otherwise noted.


<sup>1</sup>H NMR spectra were recorded on a Bruker DPX400 spectrometer using the specified solvent. Chemical shifts are expressed in parts per million (ppm,  $\delta$  units). Splitting patterns describe apparent multiplicities and are designated as s (singlet), d (doublet), t (triplet),  
 20 quartet (q), quint. (quintet), sext. (sextet) m (multiplet), br (broad), c (complex).

Low-resolution mass spectra (MS) were recorded on a JEOL JMS-AX505HA, JEOL SX-102, or a SCIEX-APIiii spectrometer; LC-MS were recorded on Waters ZQ spectrometers.

25 Preparative HPLC refers to methods where the material was purified by High Performance Liquid Chromatography on a HPLC ABZ+ 5 $\mu$ m column (10 cm x 21.2 mm i.d.) with 0.1% formic acid in water and 0.05% formic acid in acetonitrile utilising gradient elution at a flow rate of 8 mL/min and UV detection at 254 nm.

30 Mass directed auto preparation chromatography refers to methods where the material was purified by HPLC comprised of the following apparatus: Waters 600 gradient pump, Waters 2700 sample manager, Waters reagent manager, Micromass ZQ mass spectrometer, Gilson 202 - fraction collector, Gilson 202 - waste collector, Gilson 115 post-fraction UV detector. Detection is by UV and fraction collection is triggered by observation  
 35 of the programmed mass ion for the compound of interest. Software used is Micromass Masslynx version 3.5. Two columns are used: For polar (< 1.8 min analytical RT) compounds, an Optimal ODS-L is used, whose dimensions are 20mm internal diameter by 100mm in length. The stationary phase particle size is 5  $\mu$ m. For all other compounds, a Supelco LCABZ++ column is employed, whose dimensions are 20mm internal diameter by  
 40 100 mm in length. The stationary phase particle size is 5 $\mu$ m. Eluting solvents are: water + 0.1% formic acid and acetonitrile: water 95:5+0.05% formic acid; using a runtime of 14 minutes and a flow rate of 20 mL/min.




**Intermediate 1: Methyl 2-chloro-3-oxo-2,3-dihydro-2-thiophenecarboxylate**

- 5 To a solution of methyl 3-hydroxy-2-thiophenecarboxylate (5.00 g, 31.6 mmol) in chloroform (10 mL) was added 1M sulfuryl chloride in dichloromethane (34.8 mL, 34.8 mmol) dropwise over 2 minutes under a N<sub>2</sub> atmosphere. The mixture was stirred for 4 hours at room temperature and the volatiles removed under reduced pressure. The solids were recrystallized from hexane to give methyl 2-chloro-3-oxo-2,3-dihydro-2-thiophenecarboxylate (4.60 g, 76%) as white needles. <sup>1</sup>H NMR (CDCl<sub>3</sub>): δ 8.38 (d, 1 H), 6.23 (d, 1 H), 3.84 (s, 3 H); MS *m/z* 193 (M+1).

**Examples**


- 15 (1) **5-[5,6-Bis(methoxy)-1*H*-benzimidazol-1-yl]-3-([4-(hydroxymethyl)phenyl]methyl)oxy)-2-thiophenecarbonitrile**



20


- To a solution of 5-[5,6-bis(methoxy)-1*H*-benzimidazol-1-yl]-3-([4-(hydroxymethyl)phenyl]methyl)oxy)-2-thiophenecarboxamide (22.0 mg, 0.05 mmol) and pyridine (121 μL, 1.50 mmol) in dichloromethane (5 mL) cooled to approximately -10°C (ice/sodium chloride/water bath) was added trifluoroacetic anhydride (70 μL, 0.5 mmol) and the mixture stirred for 5 mins. after which the solution was allowed to warm up to room temperature over 16 h. The reaction mixture was quenched with saturated aqueous sodium hydrogen carbonate solution (1 mL) and shaken. The organic layer was recovered by passing the mixture through a PTFE phase separator, collected in a tared glass vial and concentrated to dryness by nitrogen blow down at 40 °C. The resulting crude solid was dissolved in 1:1 dimethyl sulfoxide/methanol solution (500 μL) and purified by mass directed auto preparation chromatography to afford the title compound as an off white solid (6.9 mg). MS [M+1]<sup>+</sup> 422.29; <sup>1</sup>H NMR δ<sub>H</sub> (400.13 MHz, d<sub>6</sub>-DMSO, TMS): 8.47 (s, 1 H), 7.73 (s, 1 H), 7.47 (d, *J* = 8.0 Hz, 2 H), 7.38 (d, *J* = 8.0 Hz, 2 H), 7.35 (s, 1 H), 7.23 (s, 1 H), 5.41 (s, 2 H), 5.22 (t, *J* = 5.5 Hz, 1 H), 4.52 (d, *J* = 5.5 Hz, 2 H), 3.85 (s, 3 H), 3.83 (s, 3 H).

(2) 5-[5,6-Bis(methoxy)-1*H*-benzimidazol-1-yl]-3-[[*(1R)*-1-(2-chlorophenyl)ethyl]oxy]-2-thiophenecarbonitrile



To a solution of 5-[5,6-bis(methoxy)-1*H*-benzimidazol-1-yl]-3-[[*(1R)*-1-(2-chlorophenyl)ethyl]oxy]-2-thiophenecarboxamide (31.1 mg, 0.07 mmol) and pyridine (156  $\mu$ L, 1.93 mmol) in dichloromethane (5 mL) cooled to approximately  $-10^{\circ}\text{C}$  (ice/sodium chloride/water bath) was added trifluoroacetic anhydride (408  $\mu$ L, 2.89 mmol) and the mixture stirred for 1 hr. after which the solution was poured into ice/water (10 mL). After all the ice had melted, potassium carbonate (500 mg) was added and the solution extracted with neat dichloromethane (10 mL). The organic phase was recovered by passing through a PTFE phase separator and taken down to dryness by nitrogen blow down at  $40^{\circ}\text{C}$ . The crude solid was dissolved in 1:1 dimethyl sulfoxide/methanol solution (2 mL) and purified by preparative HPLC to afford the title compound as a white solid (7.0 mg). MS  $[M+1]^+$  440.07 + 442.03 (characteristic Cl isotope pattern);  $^1\text{H}$  NMR  $\delta_{\text{H}}$  (400.13 MHz,  $\text{CD}_3\text{CN}$ , TMS): 7.96 (s, 1 H), 7.63 (dd,  $J = 1.3, 7.8$  Hz, 1 H), 7.44-7.31 (c m, 3 H), 7.26 (s, 1 H), 7.01 (s, 1 H), 6.95 (s, 1 H), (5.99, q,  $J = 6.5$  Hz, 1 H), 3.86 (s, 3 H), 3.82 (s, 3 H), 1.71 (d,  $J = 6.3$  Hz, 3 H)


(3) 5-(1*H*-Benzimidazol-1-yl)-3-[[*(2-methylphenyl)methyl*]oxy]-2-thiophenecarbonitrile



5-(1*H*-Benzimidazol-1-yl)-3-[[*(2-methylbenzyl)*oxy]-2-thiophenecarboxamide (28.5 mg, 0.08 mmol) was dissolved in pyridine (2 mL) and cooled to  $0^{\circ}\text{C}$ . Trifluoroacetic anhydride (17  $\mu$ L, 0.120 mmol) was added dropwise via syringe. The mixture was stirred for 15 min. and warmed to room temperature. After 1 h. dichloromethane (2 mL) was added followed by five drops of trifluoroacetic anhydride were added to dissolve the insoluble components of the mixture. After 14 h. the reaction was poured into dichloromethane and brine. The layers were separated and the aqueous layer was washed twice with dichloromethane. The combined organic layers were dried over magnesium sulphate, filtered, and


concentrated *in vacuo*. Purification by flash chromatography gave the title compound as a pale yellow solid (7.5 mg) MS  $[M+1]^+$  346;  $^1\text{H}$  NMR  $\delta_{\text{H}}$  (400.13 MHz,  $d_6$ -DMSO, TMS): 8.71 (s, 1H), 7.83 (s and m, 3H), 7.49-7.25 (m, 6H), 5.44 (s, 2H), 2.40 (s, 3H).

5 (4) 5-(1H-Benzimidazol-1-yl)-3-[(phenylmethyl)oxy]-2-thiophenecarbonitrile



To a solution of 5-(1H-Benzimidazol-1-yl)-3-[(phenylmethyl)oxy]-2-thiophenecarboxamide (250 mg, 0.72 mmol) in dichloromethane (5 mL) was added 2-chloro-1,3-dimethylimidazolinium chloride (125 mg, 0.74 mmol) and trifluoroacetic acid (55  $\mu\text{L}$ , 0.74 mmol). To this solution was added triethylamine (200  $\mu\text{L}$ , 1.43 mmol). The mixture was stirred for 3 h. at room temperature, then for 3 h. at reflux. The reaction mixture was cooled to room temperature and additional 2-chloro-1,3-dimethylimidazolinium chloride (125 mg, 0.74 mmol) and trifluoroacetic acid (55  $\mu\text{L}$ , 0.74 mmol) were added, followed by triethylamine (200  $\mu\text{L}$ , 1.43 mmol). The mixture was stirred for 14 h. then poured into water and extracted with dichloromethane. The organic extracts were washed with 5% aqueous hydrochloric acid, then saturated aqueous sodium hydrogen carbonate, then saturated aqueous sodium chloride solution and dried over sodium sulphate. Filtration and concentration *in vacuo*, followed by silica gel chromatography (eluting with a 20 to 50 % ethyl acetate/hexane gradient) afforded the title compound as a white solid (186 mg). MS  $[M+1]^+$  332;  $^1\text{H}$  NMR  $\delta_{\text{H}}$  (400.13 MHz,  $d_6$ -DMSO, TMS): 8.69 (s, 1H), 7.79-7.82 (m, 2H), 7.76 (s, 1H), 7.49-7.53 (m, 2H), 7.37-7.47 (m, 5H), 5.42 (s, 2H).


25 (5) 5-[5,6-Bis(methyloxy)-1H-benzimidazol-1-yl]-3-([2-(trifluoromethyl)phenyl]methyl)oxy)-2-thiophenecarbonitrile



30 To a solution of 5-(5,6-dimethoxy-1H-benzimidazol-1-yl)-3-([2-(trifluoromethyl)-benzyl]oxy)thiophene-2-carboxamide (150 mg, 0.31 mmol) in dichloromethane (2 mL) was added 2-chloro-1,3-dimethylimidazolinium chloride (120 mg, 0.71 mmol) and trifluoroacetic acid (50  $\mu\text{L}$ , 0.65 mmol). To this solution was added triethylamine (200  $\mu\text{L}$ , 1.44 mmol). The mixture was stirred for 18 h. then additional 2-chloro-1,3-dimethylimidazolinium chloride (120 mg, 0.71 mmol) and trifluoroacetic acid (50  $\mu\text{L}$ , 0.65 mmol) were added, followed by triethylamine (200  $\mu\text{L}$ , 1.44 mmol). The mixture was stirred for 4 h then poured


into water and extracted with dichloromethane. The organic extracts were washed with 5% aqueous hydrochloric acid, then saturated aqueous sodium hydrogen carbonate, then saturated aqueous sodium chloride solution and dried over sodium sulphate. Filtration and concentration followed by silica gel chromatography (eluting with a 40 to 95% ethyl acetate/hexane gradient) afforded the title compound as a yellow solid (66 mg). MS  $[M+1]^+$  459;  $^1\text{H}$  NMR  $\delta_{\text{H}}$  (400.13 MHz,  $\text{CDCl}_3$ , TMS):  $\delta$  7.90 (s, 1 H), 7.80-7.72 (m, 2 H), 7.66 (t,  $J = 7.6$  Hz, 1 H), 7.51 (t,  $J = 7.5$  Hz, 1 H), 7.31 (s, 1 H), 6.99 (s, 1 H), 6.82 (s, 1 H), 5.55 (s, 2 H), 3.96 (s, 3 H), 3.92 (s, 3 H).

10 (6) 5-[5,6-bis(methyloxy)-1*H*-benzimidazol-1-yl]-3-[[1-methyl-1*H*-1,2,3-benzotriazol-6-yl)methyl]oxy]-2-thiophenecarbonitrile



15 Prepared in a manner analogous to procedure (1) using 5-[5,6-bis(methyloxy)-1*H*-benzimidazol-1-yl]-3-[[1-methyl-1*H*-1,2,3-benzotriazol-6-yl)methyl]oxy]-2-thiophenecarboxamide (23.3 mg, 0.05 mmol). The resulting crude solid was dissolved in 1:1 dimethyl sulfoxide/methanol solution (500  $\mu\text{L}$ ) and purified by mass directed auto  
20 preparation chromatography to afford the title compound as a white solid (6.2 mg). MS  $[M+1]^+$  447.28;  $^1\text{H}$  NMR  $\delta_{\text{H}}$  (400.13 MHz,  $d_6$ -DMSO, TMS): 8.48 (s, 1 H), 8.11 (d,  $J = 8.5$  Hz, 1 H), 8.01 (s, 1 H), 7.77 (s, 1 H), 7.55 (dd,  $J = 8.5, 1.3$  Hz, 1 H), 7.35 (s, 1 H), 7.24 (s, 1 H), 5.61 (s, 2 H), 4.33 (s, 3 H), 3.83 (s, 3 H), 3.81 (s, 3 H).


25 (7) 5-(6-(Methyloxy)-5-[[2-(4-morpholinyl)ethyl]oxy]-1*H*-benzimidazol-1-yl)-3-([2-(trifluoromethyl)phenyl]methyl]oxy)-2-thiophene carbonitrile



30 Prepared in an a manner analogous to procedure (1) using 5-(6-(methyloxy)-5-[[2-(4-morpholinyl)ethyl]oxy]-1*H*-benzimidazol-1-yl)-3-([2-(trifluoromethyl)phenyl]methyl]oxy)-2-thiophenecarboxamide (17.5 mg, 0.03 mmol) The crude solid was triturated from 1:1 dimethyl sulfoxide/methanol (500  $\mu\text{L}$ ), collected by suction filtration on a PTFE phase


separator, washed well with methanol and dried under suction in air to afford the title compound as an off-white solid (12.2 mg). MS  $[M+1]^+$  559.37;  $^1\text{H}$  NMR  $\delta_{\text{H}}$  (400.13 MHz,  $d_6$ -DMSO, TMS): 8.55 (s, 1 H), 7.87-7.78 (c m, 4H), 7.67 (m, 1 H), 7.55 (s, 1 H), 7.35 (s, 1 H), 5.56 (s, 2 H), 4.39 (br. s, 2 H), 4.01 (br. s, 2 H), 3.89 (s, 3 H), 3.82-3.17 (br. c m, 8H, overlapping with residual  $\text{H}_2\text{O}$ )

(8) 5-[5,6-Bis(methyloxy)-1H-benzimidazol-1-yl]-3-[[2,5-difluorophenyl)methyl]oxy]-2-thiophenecarbonitrile




Prepared in manner analogous to procedure (1) using 5-[5,6-bis(methyloxy)-1H-benzimidazol-1-yl]-3-[[2,5-difluorophenyl)methyl]-oxy]-2-thiophenecarboxamide (22.3 mg, 0.05 mmol) and the title compound afforded by the trituration procedure described in (7) as an off-white solid (9.2 mg). MS  $[M+1]^+$  428.24;  $^1\text{H}$  NMR  $\delta_{\text{H}}$  (400.13 MHz,  $d_6$ -DMSO, TMS): 8.48 (s, 1 H), 7.79 (s, 1 H), 7.50 (c m, 1 H) 7.42-7.32, (c m, 3 H), 7.28 (s, 1 H), 5.47 (s, 2 H), 3.86 (s, 3 H), 3.83 (s, 3 H).

(9) 5-[5,6-Bis(methyloxy)-1H-benzimidazol-1-yl]-3-([2-(methylsulfonyl)phenyl)methyl]oxy)-2-thiophenecarbonitrile




Prepared in a manner analogous to procedure (1) using 5-[5,6-bis(methyloxy)-1H-benzimidazol-1-yl]-3-([2-(methylsulfonyl)phenyl)-methyl]oxy)-2-thiophenecarboxamide (24.5 mg, 0.05 mmol) The resulting crude solid was dissolved in 1:1 dimethyl sulfoxide/methanol solution (500  $\mu\text{L}$ ) and purified by mass directed auto preparation chromatography to afford the title compound as an off-white solid (4.6 mg). MS  $[M+1]^+$  470.25;  $^1\text{H}$  NMR  $\delta_{\text{H}}$  (400.13 MHz,  $d_6$ -DMSO, TMS): 8.50 (s, 1 H), 8.06 (d,  $J = 7.8$ , 1 H), 7.86-7.83 (c m, 2 H), 7.76-7.72 (cm, 2 H), 7.36 (s, 1 H), 7.28 (s, 1 H), 5.82 (s, 2 H), 3.86 (s, 3 H), 3.83 (s, 3 H), 3.35 (s, 3 H)

(10) 5-(5-Chloro-1H-benzimidazol-1-yl)-3-[(phenylmethyl)oxy]-2-thiophenecarbonitrile




Prepared in a manner analogous to procedure (4) using 5-(5-Chloro-1*H*-benzimidazol-1-yl)-3-[(phenylmethyl)oxy]-2-thiophenecarboxamide (105 mg, 0.270 mmol). Silica gel chromatography (eluting with a 10 to 40% solution of ethyl acetate in hexane) afforded the title compound as an off-white solid (71 mg) MS  $[M+1]^+$  366;  $^1\text{H}$  NMR  $\delta_{\text{H}}$  (400.13 MHz,  $d_6$ -DMSO, TMS): 8.75 (s, 1 H), 7.91 (d,  $J$  = 1.8 Hz, 1 H), 7.79-7.82 (m, 1 H), 7.70 (s, 1 H), 7.39-7.52 (m, 6 H), 5.41 (s, 2 H).

(11) 5-[5,6-bis(Methyloxy)-1*H*-benzimidazol-1-yl]-3-[[2,6-difluorophenyl)methyl]oxy]-2-thiophenecarbonitrile



Prepared in a manner analogous to procedure (1) using 5-[5,6-bis(methyloxy)-1*H*-benzimidazol-1-yl]-3-[[2,6-difluorophenyl)-methyl]-oxy]-2-thiophenecarboxamide (22.4 mg, 0.05 mmol) and the title compound afforded by the trituration procedure described in (7) as an off-white solid (9.9 mg). MS  $[M+1]^+$  428.24;  $^1\text{H}$  NMR  $\delta_{\text{H}}$  (400.13 MHz,  $d_6$ -DMSO, TMS): 8.48 (s, 1 H), 7.81 (s, 1 H), 7.60 (tt,  $J$  = 8.5, 6.8 Hz, 1 H), 7.36 (s, 1 H), 7.33 (s, 1 H), 7.25 (t,  $J$  = 8.0 Hz, 2 H), 5.49 (s, 2 H), 3.88 (s, 3 H), 3.83 (s, 3 H).


(12) 5-[5-[(3-Hydroxypropyl)oxy]-6-(methyloxy)-1*H*-benzimidazol-1-yl]-3-[[2-(trifluoromethyl)phenyl]methyl]oxy]-2-thiophenecarbonitrile



Prepared in a manner analogous to procedure (1) using 5-[5-[(3-hydroxypropyl)oxy]-6-(methyloxy)-1*H*-benzimidazol-1-yl]-3-[[2-(trifluoromethyl)phenyl]methyl]oxy)-2-thiophenecarboxamide (26.1 mg, 0.05 mmol) The resulting crude solid was dissolved in


1:1 dimethyl sulfoxide/methanol solution (500  $\mu$ L) and purified by mass directed auto preparation chromatography to afford the title compound as an off-white solid (5.4 mg). MS  $[M+1]^+$  504.27;  $^1\text{H}$  NMR  $\delta_{\text{H}}$  (400.13 MHz,  $d_6$ -DMSO, TMS): 8.49 (s, 1 H), 7.86-7.78 (c m, 4H), 7.66 (m, 1 H), 7.35 (s, 1 H), 7.28 (s, 1 H), 5.56 (s, 2 H), 4.55 (t,  $J$  = 5.0 Hz, 1 H), 4.09 (t,  $J$  = 6.3 Hz, 2 H), 3.86 (s, 3 H), 3.58 (dd,  $J$  = 6.0, 5.5 Hz, 2 H), 1.90 (quint.,  $J$  = 6.3 Hz, 2 H).

(13) 5-[5,6-Bis(methoxy)-1*H*-benzimidazol-1-yl]-3-[[3-bromo-4-pyridinyl)methyl]oxy]-2-thiophenecarbonitrile




Prepared in a manner analogous to procedure (1) using 5-[5,6-bis(methoxy)-1*H*-benzimidazol-1-yl]-3-[[3-bromo-4-pyridinyl)methyl]oxy]-2-thiophenecarboxamide (24.4 mg, 0.05 mol) and the title compound afforded by the trituration procedure described in (7) as an off-white solid (7.6 mg). MS  $[M+1]^+$  471.19 + 473.19 (characteristic Br isotope pattern);  $^1\text{H}$  NMR  $\delta_{\text{H}}$  (400.13 MHz,  $d_6$ -DMSO, TMS): 8.81 (s, 1 H), 8.66 (d,  $J$  = 5.0 Hz, 1 H), 8.49 (s, 1 H), 7.76 (s, 1 H), 7.61 (d,  $J$  = 4.8 Hz, 1 H), 7.35 (s, 1 H), 7.26 (s, 1 H), 5.52 (s, 2 H), 3.84 (s, 3 H), 3.83 (s, 3 H).

(14) 5-[5,6-Bis(methoxy)-1*H*-benzimidazol-1-yl]-3-[(cyclohexylmethyl)oxy]-2-thiophenecarbonitrile




Prepared in manner analogous to procedure (1) using 5-[5,6-bis(methoxy)-1*H*-benzimidazol-1-yl]-3-[(cyclohexylmethyl)oxy]-2-thiophenecarboxamide (20.9 mg, 0.05 mmol). The resulting crude solid was dissolved in 1:1 dimethyl sulfoxide/methanol solution (500  $\mu$ L) and purified by mass directed auto preparation chromatography to afford the title compound as an off white solid (5.8 mg). MS  $[M+1]^+$  398.35;  $^1\text{H}$  NMR  $\delta_{\text{H}}$  (400.13 MHz,  $d_6$ -DMSO, TMS): 8.48 (s, 1 H), 7.63 (s, 1 H), 7.35 (s, 1 H), 7.28 (s, 1 H), 4.14 (d,  $J$  = 6.0 Hz, 2 H), 3.86 (s, 3 H), 3.83 (s, 3 H), 1.82-1.61 (br. c m, 6H), 1.32-1.01 (br. c m, 5H).

(15) 1-[5-cyano-4-([2-(trifluoromethyl)phenyl)methyl]oxy)-2-thienyl]-*N*-[2-(4-morpholinyl)ethyl]-1*H*-benzimidazole-5-carboxamide



Prepared in an analogous manner to procedure (1) using 1-[5-(aminocarbonyl)-4-({[2-(trifluoromethyl)phenyl]methyl}oxy)-2-thienyl]-N-[2-(4-morpholinyl)ethyl]-1H-benzimidazole-5-carboxamide (28.7 mg, 0.05 mmol). The resulting crude solid was dissolved in 1:1 dimethyl sulfoxide/methanol solution (500  $\mu$ L) and purified by mass directed auto preparation chromatography to afford the title compound as an off white solid (4.4 mg). MS  $[M+1]^+$  556.36;  $^1\text{H}$  NMR  $\delta_{\text{H}}$  (400.13 MHz,  $d_6$ -DMSO, TMS): 8.80 (s, 1 H), 8.55 (t,  $J$  = 5.5 Hz, 1 H), 8.33 (s, 1 H), 7.97, (m, 1 H), 7.90-7.77 (c m, 5H), 7.67 (t,  $J$  = 7.5, 1 H), 5.56 (s, 2 H), 3.58 (t,  $J$  = 4.5 Hz, 4H), 3.43 (q, 6.8 Hz, 2 H), 2.43 (br. s, 4H), other signal (2 H) too broad to observe.


(16) 5-[5,6-Bis(methyloxy)-1H-benzimidazol-1-yl]-3-({(1R)-1-[2-(trifluoromethyl)phenyl]ethyl}oxy)-2-thiophenecarbonitrile



Prepared in an analogous manner to procedure (1) using 5-[5,6-bis(methyloxy)-1H-benzimidazol-1-yl]-3-({(1R)-1-[2-(trifluoromethyl)phenyl]ethyl}oxy)-2-thiophenecarboxamide (24.4 mg, 0.05 mmol). The crude solid was partitioned between dichloromethane (10 mL) and 2M hydrochloric acid solution (5 mL), the organic phase was recovered by passing through a PTFE phase separator and concentrated to dryness by nitrogen blowdown at 40  $^{\circ}\text{C}$  to afford the title compound as an off-white solid (8.2 mg). MS  $[M+1]^+$  474.31;  $^1\text{H}$  NMR  $\delta_{\text{H}}$  (400.13 MHz,  $d_6$ -DMSO, TMS): 8.49 (s, 1 H), 7.89-7.75 (c m, 3 H), 7.58 (m, 1 H), 7.41 (s, 1 H), 7.34 (s, 1 H), 7.13 (s, 1 H), 6.04 (q,  $J$  = 5.8 Hz, 1 H), 3.82 (s, 3 H), 3.81 (s, 3 H), 1.71 (d,  $J$  = 6.0 Hz, 3 H).


(17) 5-[5,6-bis(methyloxy)-1H-benzimidazol-1-yl]-3-[(2-phenylethyl)oxy]-2-thiophenecarbonitrile






Prepared in an analogous manner to procedure (1) using 5-[5,6-bis(methoxy)-1H-benzimidazol-1-yl]-3-[(2-phenylethyl)oxy]-2-thiophenecarbonitrile (21.0 mg, 0.05 mmol). The crude solid was dissolved in 1:1 dimethyl sulfoxide/methanol solution (0.5 mL) and purified by preparative HPLC to afford the title compound as a white solid (5.1 mg). MS [M+1]<sup>+</sup> 406.14; <sup>1</sup>H NMR  $\delta_{\text{H}}$  (400.13 MHz, *d*<sub>6</sub>-DMSO, TMS): 8.47 (s, 1 H), 7.65 (s, 1 H), 7.37-7.23 (cm, 7H), 4.54 (t, *J* = 6.8 Hz, 2 H), 3.85 (s, 3 H), 3.82 (s, 3 H), 3.10 (t, *J* = 6.8 Hz, 2 H).

- 10 (18) 5-[5,6-bis(methoxy)-1H-benzimidazol-1-yl]-3-[(2-[(trifluoromethyl)oxy]phenyl)methyl]oxy]-2-thiophenecarbonitrile




Prepared in an analogous manner to procedure (1) using 5-[5,6-bis(methoxy)-1H-benzimidazol-1-yl]-3-[(2-[(trifluoromethyl)oxy]phenyl)methyl]oxy]-2-thiophenecarbonitrile (24.6 mg, 0.05 mmol). The crude solid was dissolved in 1:1 dimethyl sulfoxide/methanol solution (0.5 mL) and purified by preparative HPLC to afford the title compound as a white solid (3.2 mg). MS [M+1]<sup>+</sup> 476.09; <sup>1</sup>H NMR  $\delta_{\text{H}}$  (400.13 MHz, *d*<sub>6</sub>-DMSO, TMS): 8.49 (s, 1 H), 7.80 (s, 1 H), 7.75 (dd, *J* = 7.5, 1.3 Hz, 1 H), 7.58 (dd, *J* = 7.8, 1.5 Hz, 1 H), 7.51-7.47 (cm, 2 H), 7.36 (s, 1 H), 7.28 (s, 1 H), 5.48 (s, 2 H), 3.86 (s, 3 H), 3.83 (s, 3 H).

- (19) 5-[5,6-bis(methoxy)-1H-benzimidazol-1-yl]-3-[(2-(methoxy)phenyl)methyl]oxy]-2-thiophenecarbonitrile




Prepared in an analogous manner to procedure (1) using 5-[5,6-bis(methoxy)-1*H*-benzimidazol-1-yl]-3-([2-(methoxy)phenyl]-methyl)oxy)-2-thiophenecarbonitrile (22 mg, 0.05 mmol). The crude solid was dissolved in 1:1 dimethyl sulfoxide/methanol solution (0.5 mL) and purified by preparative HPLC to afford the title compound as a white solid (4.9 mg). MS  $[M+1]^+$  422.16;  $^1\text{H}$  NMR  $\delta_{\text{H}}$  (400.13 MHz,  $d_6$ -DMSO, TMS): 8.50 (s, 1 H), 7.74 (s, 1 H), 7.48 (d,  $J = 7.5$  Hz, 1 H), 7.40 (dt,  $J = 1.5, 8.3$  Hz, 1 H), 7.63 (s, 1 H), 7.28 (s, 1 H), 7.10 (d,  $J = 8.3$ , 1 H), 7.01 (t,  $J = 7.3$  Hz), 5.39 (s, 2 H), 3.85 (s, 3 H), 3.84 (s, 3 H), 3.83 (s, 3 H).

10 (20) 5-[5,6-bis(methoxy)-1*H*-benzimidazol-1-yl]-3-([2-(4-morpholinyl)ethyl]oxy)-2-thiophenecarbonitrile



15 Prepared in an analogous manner to procedure (1) using 5-[5,6-bis(methoxy)-1*H*-benzimidazol-1-yl]-3-([2-(4-morpholinyl)-ethyl]oxy)-2-thiophenecarboxamide (21.5 mg, 0.05 mmol). The crude solid was subject to the partitioning procedure described in example (16) to afford the title compound as a colourless oil (5.2 mg). MS  $[M+1]^+$  415.29;  $^1\text{H}$  NMR  $\delta_{\text{H}}$  (400.13 MHz,  $d_6$ -DMSO, TMS): 8.47 (s, 1 H), 7.65 (s, 1 H), 7.35 (s, 1 H), 7.29 (2, 1 H), 4.44 (t,  $J = 5.5$  Hz, 2 H), 3.86 (s, 3 H), 3.83 (s, 3 H), 3.57 (t,  $J = 4.5$  Hz, 4H), 2.74 (t,  $J = 5.5$  Hz, 2 H), 2.50 (4H multiplet obscured by DMSO signal)


20 (21) 5-[5,6-bis(methoxy)-1*H*-benzimidazol-1-yl]-3-([2-(2-oxo-1-pyrrolidinyl)ethyl]oxy)-2-thiophenecarbonitrile



25 To a solution of 5-[5,6-bis(methoxy)-1*H*-benzimidazol-1-yl]-3-([2-(2-oxo-1-pyrrolidinyl)-ethyl]oxy)-2-thiophenecarboxamide (15.2 mg, 0.035 mmol) and pyridine (121  $\mu\text{L}$ , 1.50 mmol) in dichloromethane (5 mL) cooled to approximately  $-10^\circ\text{C}$  (ice/sodium chloride/water bath) was added trifluoroacetic anhydride (70  $\mu\text{L}$ , 0.5 mmol) and the mixture  
30 stirred for 5 mins. after which the solution was allowed to warm up to room temperature


over 16 h. The reaction was quenched with 2M aqueous sodium carbonate solution (2 mL) and shaken. The organic layer was recovered by passing the mixture through a PTFE phase separator, collected in a tared glass vial and concentrated to dryness by nitrogen blow down at 40 °C. The resultant crude solid was ultrasonicated in methanol (10 mL) and loaded onto a pre-equilibrated (methanol, 9 mL) aminopropyl functionalised silica cartridge (Isolute, 500 mg) and three column volumes of methanol (9 mL) were allowed to percolate under gravity and collected in tared glass vial. Concentration in a vacuum centrifuge at 50 °C afforded the title compound as an off-white solid (4.9 mg). MS  $[M+1]^+$  413.14;  $^1\text{H}$  NMR  $\delta_{\text{H}}$  (400.13 MHz,  $d_6$ -DMSO, TMS): 8.47 (s, 1 H), 7.65 (s, 1 H), 7.35 (s, 1 H), 7.30 (s, 1 H), 4.44 (t,  $J$  = 5.3 Hz, 2 H), 3.86 (s, 3 H), 3.83 (s, 3 H), 3.59 (t,  $J$  = 5.3 Hz, 2 H), 3.50 (t,  $J$  = 7.0 Hz, 2 H), 2.23 (t,  $J$  = 7.8 Hz, 2 H), 1.94 (quint.,  $J$  = 7.5 Hz (average), 2 H).

(22) 5-[5,6-bis(methyloxy)-1*H*-benzimidazol-1-yl]-3-[(tetrahydro-2-furanylmethyl)oxy]-2-thiophenecarbonitrile



Prepared and isolated in a manner analogous to procedure (21) using 5-[5,6-bis(methyloxy)-1*H*-benzimidazol-1-yl]-3-[(tetrahydro-2-furanylmethyl)oxy]-2-thiophenecarboxamide (14.1 mg, 0.035 mmol). The title compound was isolated as an off-white solid (3.3 mg). MS  $[M+1]^+$  386.12;  $^1\text{H}$  NMR  $\delta_{\text{H}}$  (400.13 MHz,  $d_6$ -DMSO, TMS): 8.47 (s, 1 H), 7.63 (s, 1 H), 7.35 (s, 1 H), 7.28 (s, 1 H), 4.39-4.35 (c m, 1 H), 4.30-4.17 (c m, 2 H), 3.86 (s, 3 H), 3.83 (s, 3 H), 3.79 (t,  $J$  = 6.8 Hz, 1 H), 3.71 (quint.,  $J$  = 7.3 Hz (average), 1 H), 2.07-1.79 (c m, 3 H), 1.72-1.64 (c m, 1 H).


(23) 5-[5,6-bis(methyloxy)-1*H*-benzimidazol-1-yl]-3-[(tetrahydro-2H-pyran-2-ylmethyl)oxy]-2-thiophenecarbonitrile



Prepared and isolated in a manner analogous to procedure (21) using 5-[5,6-bis(methyloxy)-1*H*-benzimidazol-1-yl]-3-[(tetrahydro-2H-pyran-2-ylmethyl)oxy]-2-thiophene-carboxamide (14.6 mg, 0.035 mmol). The title compound was isolated as an off-white solid (3.3 mg). MS  $[M+1]^+$  400.12;  $^1\text{H}$  NMR  $\delta_{\text{H}}$  (400.13 MHz,  $d_6$ -DMSO, TMS): 8.47 (s, 1 H), 7.62 (s, 1 H), 7.35 (s, 1 H), 7.28 (s, 1 H), 4.29 (d,  $J$  = 5.0 Hz, 2 H), 3.91 (m, 1 H), 3.86


(s, 3 H), 3.83 (s, 3 H), 3.70-3.66 (cm, 1 H), 3.44-3.37 (cm, 1 H), 1.96-1.81 (cm, 1 H), 1.67-1.43 (cm, 4H), 1.37-1.27 (cm, 1 H).

(24) 5-[5,6-bis(methyloxy)-1*H*-benzimidazol-1-yl]-3-[[2-(phenyloxy)ethyl]oxy]-2-thiophenecarbonitrile




Prepared and isolated in a manner analogous to procedure (21) using 5-[5,6-bis(methyloxy)-1*H*-benzimidazol-1-yl]-3-[[2-(phenyl-oxy)ethyl]oxy]-2-thiophenecarboxamide (15.7 mg, 0.035 mmol). The title compound was isolated as an off-white solid (3.3 mg). MS  $[M+1]^+$  422.12;  $^1\text{H}$  NMR  $\delta_{\text{H}}$  (400.13 MHz,  $d_6$ -DMSO, TMS): 8.48 (s, 1 H), 7.70 (s, 1 H), 7.35 (s, 1 H), 7.33-7.29 (m, 3 H), 7.00-6.95 (m, 3 H), 4.70 (m, 2 H), 4.37 (m, 2 H), 3.85 (s, 3 H), 3.83 (s, 3 H).

(25) 5-[5,6-bis(methyloxy)-1*H*-benzimidazol-1-yl]-3-[[1*S*]-1-(2-chlorophenyl)butyl]oxy]-2-thiophenecarbonitrile




Prepared and isolated in a manner analogous to procedure (21) using 5-[5,6-bis(methyloxy)-1*H*-benzimidazol-1-yl]-3-[[1*S*]-1-(2-chlorophenyl)butyl]oxy]-2-thiophenecarboxamide (17.4 mg, 0.035 mmol). The title compound was isolated as an off-white solid (5.7 mg). MS  $[M+1]^+$  468.09 + 470.08 (characteristic chlorine isotope pattern);  $^1\text{H}$  NMR  $\delta_{\text{H}}$  (400.13 MHz,  $d_6$ -DMSO, TMS): 8.41 (s, 1 H), 7.57 (dd,  $J = 1.0, 7.53$  Hz, 1 H), 7.48 (d,  $J = 7.8$  Hz, 1 H), 7.44 (t,  $J = 6.8$  Hz, 1 H), 7.38-7.32 (cm, 3 H), 7.08 (s, 1 H), 5.90 (m, 1 H), 3.81 (s, 1 H), 3.80 (s, 3 H), 2.07-1.86 (cm, 2 H), 1.56-1.37 (m, 2 H), 0.96 (t,  $J = 7.3$  Hz, 3 H).

(26) 5-[5,6-bis(methyloxy)-1*H*-benzimidazol-1-yl]-3-[(3-thienylmethyl)oxy]-2-thiophenecarbonitrile




Prepared and isolated in a manner analogous to procedure (21) using 5-[5,6-bis(methoxy)-1*H*-benzimidazol-1-yl]-3-[(3-thienyl-methyl)oxy]-2-thiophenecarboxamide (14.9 mg, 0.035 mmol). The title compound was isolated as an off-white solid (mg). MS [M+1]<sup>+</sup> 398.08; <sup>1</sup>H NMR δ<sub>H</sub> (400.13 MHz, d<sub>6</sub>-DMSO, TMS): 8.47 (s, 1 H), 7.74 (overlapping signals, 2 H), 7.62 (dd, *J* = 3.0, 4.8 Hz, 1 H), 7.36 (s, 1 H), 7.24 (overlapping signals, 2 H), 5.42 (s, 2 H), 3.86 (s, 3 H), 3.83 (s, 3 H).

(27) 5-[5,6-bis(methoxy)-1*H*-benzimidazol-1-yl]-3-[(2-thienylmethyl)oxy]-2-thiophenecarbonitrile



Prepared and isolated in a manner analogous to procedure (21) using 5-[5,6-bis(methoxy)-1*H*-benzimidazol-1-yl]-3-[(2-thienylmethyl)oxy]-2-thiophenecarboxamide (14.5 mg, 0.035 mmol). The title compound was isolated as a clear oil (7.0 mg). MS [M+1]<sup>+</sup> 398.08; <sup>1</sup>H NMR δ<sub>H</sub> (400.13 MHz, d<sub>6</sub>-DMSO, TMS): 8.47 (s, 1 H), 7.79 (s, 1 H), 7.66 (d, *J* = 5.3 Hz, 1 H), 7.36 (overlapping signals, 2 H), 7.25 (s, 1 H), 7.08 (dd, *J* = 3.5, 4.8 Hz, 1 H), 5.63 (s, 2 H), 3.87 (s, 3 H), 3.83 (s, 3 H).

(28) 5-[5,6-bis(methoxy)-1*H*-benzimidazol-1-yl]-3-[(1*R*)-1-methylpropyl]oxy]-2-thiophenecarbonitrile



Prepared and isolated in a manner analogous to procedure (21) using 5-[5,6-bis(methoxy)-1*H*-benzimidazol-1-yl]-3-[(1*R*)-1-methylpropyl]oxy]-2-thiophenecarboxamide (13.6 mg, 0.035 mmol). The title compound was isolated as an off-white solid (5.3 mg). MS [M+1]<sup>+</sup> 358.11; <sup>1</sup>H NMR δ<sub>H</sub> (400.13 MHz, d<sub>6</sub>-DMSO, TMS): 8.49 (s,

1 H), 7.64 (s, 1 H), 7.35 (s, 1 H), 7.28 (s, 1 H), 4.66 (sext.,  $J = 6.0$  Hz, 1 H), 3.86 (s, 3 H), 3.83 (s, 3 H), 1.71 (cm, 2 H), 1.34 (d,  $J = 6.3$  Hz, 3 H), 0.96 (t,  $J = 7.3$  Hz, 3 H).

## 5 Biological data

### IKK3 assay

Recombinant human IKK3 was expressed in baculovirus as a FLAG tagged fusion protein, and its activity was assessed using a time-resolved fluorescence resonance energy transfer (TR-FRET) assay. Briefly, IKK3 (15 nM final) diluted in assay buffer (50 mM HEPES, 10 mM  $MgCl_2$ , 1 mM CHAPS pH 7.4 with 1 mM DTT and 0.01% w/v BSA) was added to wells containing various concentrations of compound or DMSO vehicle (3% final). The reaction was initiated by the addition of GST-I $\kappa$ B $\alpha$  substrate (25 nM final)/ATP (2  $\mu$ M final), in a total volume of 30  $\mu$ l. The reaction was incubated for 15 minutes at room temperature, then terminated by the addition of 15  $\mu$ l of 50 mM EDTA. Detection reagent (15  $\mu$ l) in buffer (100 mM HEPES pH 7.4, 150 mM NaCl and 0.1% w/v BSA) containing antiphosphoserine-I $\kappa$ B $\alpha$ -32/36 monoclonal antibody 12C2 (Cell Signalling Technology, Beverly Massachusetts, USA) labelled with W-1024 europium chelate (Wallac OY, Turku, Finland), and an APC-labelled anti-GST antibody (Prozyme, San Leandro, California, USA) was added and the reaction was further incubated for 45 minutes at room temperature. The degree of phosphorylation of GST-I $\kappa$ B $\alpha$  was measured using a Packard Discovery plate reader (Perkin-Elmer Life Sciences, Pangbourne, UK) as a ratio of specific 665 nm energy transfer signal to reference europium 620 nm signal.

25 The results are shown in the table below:

Example	IKK3 Activity
16	*
13	*
28	*
22	*
2	*
20	*
10	*
19	*
4	*
3	**
21	**
24	**
15	**
23	**
1	**

25	**
17	**
18	**
27	**
5	**
6	**
14	**
8	**
26	**
11	**
7	**
9	**
12	**


\* =  $pIC_{50}$  of 5.0 - 6.0;    \*\* =  $pIC_{50}$  of >6.0;

- 5 The application of which this description and claims forms part may be used as a basis for priority in respect of any subsequent application. The claims of such subsequent application may be directed to any novel feature or combination of features described herein. This may take the form of product, composition, process or use claims and may include, by way of example and without limitation, one or more of the following claims.

## CLAIMS

We claim:

- 5 1. A compound of Formula (I):



wherein

10

n is 0, 1, 2, 3, or 4;

Each R<sup>1</sup> which may be the same or different, independently represents H, halogen or a group (X)<sub>a</sub>(Y)<sub>b</sub>Z;

X represents -O- or -CONH-;

15

a is 0 or 1;

Y represents -C<sub>1-6</sub> alkylene-

b is 0 or 1;

Z represents hydroxy, C<sub>1-6</sub> alkyl, C<sub>1-6</sub> haloalkyl, C<sub>5-7</sub> heterocyclyl, C<sub>1-6</sub> alkoxyalkyl, C<sub>1-6</sub> haloalkoxyalkyl;

20

R<sup>2</sup> represents a group -(X<sup>1</sup>)<sub>c</sub>(Y<sup>1</sup>)<sub>d</sub>Z<sup>1</sup>

Wherein X<sup>1</sup> represents -C<sub>1-12</sub> alkylene-;

c is 0 or 1;

25

Y<sup>1</sup> represents -O-;

d is 0 or 1;

Z<sup>1</sup> represents H, aryl or heteroaryl each of which contains 5 - 14 ring atoms, C<sub>5-7</sub> heterocyclyl, C<sub>5-7</sub> cycloalkyl, C<sub>5-7</sub> cycloalkenyl, each of which may be optionally substituted by one or more substituents independently selected from C<sub>1-6</sub> alkyl, C<sub>1-6</sub> haloalkyl, halogen,

30

C<sub>1-6</sub> alkoxy, C<sub>1-6</sub> haloalkoxy, SO<sub>2</sub>R<sup>3</sup>, C<sub>1-6</sub> hydroxyalkyl.


R<sup>3</sup> represents H or C<sub>1-6</sub> alkyl;

or pharmaceutically acceptable salts, solvates or physiologically functional derivatives thereof with the proviso that the compound is not:


35

5-(1*H*-benzimidazol-1-yl)-3-[(2-methylphenyl)oxy]thiophene-2-carbonitrile





or 5-(5,6-bismethoxy-1*H*-benzimidazol-1-yl)-3-((2-trifluoromethyl)phenyl)methoxythiophene-2-carbonitrile



5

2. A compound according to claim 1, selected from:

5-[5,6-bis(methoxy)-1 <i>H</i> -benzimidazol-1-yl]-3-([4-(hydroxymethyl)phenyl]methyl)oxy)-2-thiophenecarbonitrile
5-[5,6-bis(methoxy)-1 <i>H</i> -benzimidazol-1-yl]-3-[(1 <i>R</i> )-1-(2-chlorophenyl)ethyl]oxy)-2-thiophenecarbonitrile
5-(1 <i>H</i> -benzimidazol-1-yl)-3-[(phenylmethyl)oxy]-2-thiophenecarbonitrile
5-[5,6-bis(methoxy)-1 <i>H</i> -benzimidazol-1-yl]-3-[(1-methyl-1 <i>H</i> -1,2,3-benzotriazol-6-yl)methyl]oxy)-2-thiophenecarbonitrile)
5-(6-(methoxy)-5-[[2-(4-morpholinyl)ethyl]oxy]-1 <i>H</i> -benzimidazol-1-yl)-3-([2-(trifluoromethyl)phenyl]methyl)oxy)-2-thiophenecarbonitrile
5-[5,6-bis(methoxy)-1 <i>H</i> -benzimidazol-1-yl]-3-[(2,5-difluorophenyl)methyl]oxy)-2-thiophenecarbonitrile
5-[5,6-bis(methoxy)-1 <i>H</i> -benzimidazol-1-yl]-3-([2-(methylsulfonyl)phenyl]methyl)oxy)-2-thiophenecarbonitrile
5-(5-chloro-1 <i>H</i> -benzimidazol-1-yl)-3-[(phenylmethyl)oxy]-2-thiophenecarbonitrile
5-[5,6-bis(methoxy)-1 <i>H</i> -benzimidazol-1-yl]-3-[(2,6-difluorophenyl)methyl]oxy)-2-thiophenecarbonitrile
5-[5-[(3-hydroxypropyl)oxy]-6-(methoxy)-1 <i>H</i> -benzimidazol-1-yl]-3-([2-(trifluoromethyl)phenyl]methyl)oxy)-2-thiophene-carbonitrile
5-[5,6-bis(methoxy)-1 <i>H</i> -benzimidazol-1-yl]-3-[(3-bromo-4-pyridinyl)methyl]oxy)-2-thiophenecarbonitrile
5-[5,6-bis(methoxy)-1 <i>H</i> -benzimidazol-1-yl]-3-[(cyclohexylmethyl)oxy]-2-

thiophenecarbonitrile
1-[5-cyano-4-({[2-(trifluoromethyl)phenyl]methyl}oxy)-2-thienyl]-N-[2-(4-morpholinyl)ethyl]-1H-benzimidazole-5-carboxamide
5-[5,6-bis(methyloxy)-1H-benzimidazol-1-yl]-3-({(1R)-1-[2-(trifluoromethyl)phenyl]ethyl}oxy)-2-thiophenecarbonitrile
5-[5,6-bis(methyloxy)-1H-benzimidazol-1-yl]-3-[(2-phenylethyl)oxy]-2-thiophenecarbonitrile
5-[5,6-bis(methyloxy)-1H-benzimidazol-1-yl]-3-[(2-((trifluoromethyl)oxy)phenyl)methyl]oxy]-2-thiophenecarbonitrile
5-[5,6-bis(methyloxy)-1H-benzimidazol-1-yl]-3-({[2-(methyloxy)phenyl]methyl}oxy)-2-thiophenecarbonitrile
5-[5,6-bis(methyloxy)-1H-benzimidazol-1-yl]-3-{[2-(4-morpholinyl)ethyl]oxy}-2-thiophenecarbonitrile
5-[5,6-bis(methyloxy)-1H-benzimidazol-1-yl]-3-{[2-(2-oxo-1-pyrrolidinyl)ethyl]oxy}-2-thiophenecarbonitrile
5-[5,6-bis(methyloxy)-1H-benzimidazol-1-yl]-3-[(tetrahydro-2-furanylmethyl)oxy]-2-thiophenecarbonitrile
5-[5,6-bis(methyloxy)-1H-benzimidazol-1-yl]-3-[(tetrahydro-2H-pyran-2-ylmethyl)oxy]-2-thiophenecarbonitrile
5-[5,6-bis(methyloxy)-1H-benzimidazol-1-yl]-3-[[2-(phenyloxy)ethyl]oxy]-2-thiophenecarbonitrile
5-[5,6-bis(methyloxy)-1H-benzimidazol-1-yl]-3-[[1-(2-chlorophenyl)butyl]oxy]-2-thiophenecarbonitrile
5-[5,6-bis(methyloxy)-1H-benzimidazol-1-yl]-3-[(3-thienylmethyl)oxy]-2-thiophenecarbonitrile
5-[5,6-bis(methyloxy)-1H-benzimidazol-1-yl]-3-[(2-thienylmethyl)oxy]-2-thiophenecarbonitrile
5-[5,6-bis(methyloxy)-1H-benzimidazol-1-yl]-3-[[1-(1-methylpropyl)oxy]-2-thiophenecarbonitrile

or a salt, solvate, or physiologically functional derivative thereof.

3. A pharmaceutical composition, comprising: a therapeutically effective amount of a compound as claimed in any one of claims 1 or 2, or a salt, solvate, or a physiologically functional derivative thereof and one or more of pharmaceutically acceptable carriers, diluents and excipients.
4. A compound as claimed in any of claims 1 or 2, or a salt, solvate, or a physiologically functional derivative thereof for use in therapy.
5. A method of treating a disorder in a mammal, said disorder being mediated by at least one of inappropriate IKK3 activity, comprising: administering to said mammal a

therapeutically effective amount of a compound of Formula (Ia), or a salt, or a physiologically functional derivative thereof.

6. A method according to claim 5 wherein the disorder mediated by inappropriate IKK3 activity is inflammatory bowel disease, asthma and COPD (chronic obstructive pulmonary disease); osteoarthritis, osteoporosis and fibrotic diseases; dermatosis, including psoriasis, atopic dermatitis and ultraviolet radiation (UV)-induced skin damage; autoimmune diseases including systemic lupus erythematosus, multiple sclerosis, psoriatic arthritis, ankylosing spondylitis, tissue and organ rejection, Alzheimer's disease, stroke, atherosclerosis, restenosis, diabetes, glomerulonephritis, cancer, including Hodgkins disease, cachexia, inflammation associated with infection and certain viral infections, including acquired immune deficiency syndrome (AIDS), adult respiratory distress syndrome, and Ataxia Telangiectasia.
7. A method according to claim 6 or 7 wherein the compound of Formula (Ia) is selected from:

5-[5,6-bis(methoxy)-1H-benzimidazol-1-yl]-3-({[4-(hydroxymethyl)phenyl]methyl}oxy)-2-thiophenecarbonitrile
5-[5,6-bis(methoxy)-1H-benzimidazol-1-yl]-3-[(1R)-1-(2-chlorophenyl)ethyl]oxy)-2-thiophenecarbonitrile
5-(1H-benzimidazol-1-yl)-3-({[2-methylphenyl]methyl}oxy)-2-thiophenecarbonitrile
5-(1H-benzimidazol-1-yl)-3-[(phenylmethyl)oxy]-2-thiophenecarbonitrile
5-[5,6-bis(methoxy)-1H-benzimidazol-1-yl]-3-({[1-methyl-1H-1,2,3-benzotriazol-6-yl]methyl}oxy)-2-thiophenecarbonitrile)
5-(6-(methoxy)-5-{{2-(4-morpholinyl)ethyl}oxy}-1H-benzimidazol-1-yl)-3-({[2-(trifluoromethyl)phenyl]methyl}oxy)-2-thiophenecarbonitrile
5-[5,6-bis(methoxy)-1H-benzimidazol-1-yl]-3-({[2,5-difluorophenyl]methyl}oxy)-2-thiophenecarbonitrile
5-[5,6-bis(methoxy)-1H-benzimidazol-1-yl]-3-({[2-(trifluoromethyl)phenyl]methyl}oxy)-2-thiophenecarbonitrile
5-[5,6-bis(methoxy)-1H-benzimidazol-1-yl]-3-({[2-(methylsulfonyl)phenyl]methyl}oxy)-2-thiophenecarbonitrile
5-(5-chloro-1H-benzimidazol-1-yl)-3-[(phenylmethyl)oxy]-2-thiophenecarbonitrile
5-[5,6-bis(methoxy)-1H-benzimidazol-1-yl]-3-({[2,6-difluorophenyl]methyl}oxy)-2-thiophenecarbonitrile
5-[5-[(3-hydroxypropyl)oxy]-6-(methoxy)-1H-benzimidazol-1-yl]-3-({[2-(trifluoromethyl)phenyl]methyl}oxy)-2-thiophene-carbonitrile
5-[5,6-bis(methoxy)-1H-benzimidazol-1-yl]-3-({[3-bromo-4-pyridinyl]methyl}oxy)-2-thiophenecarbonitrile
5-[5,6-bis(methoxy)-1H-benzimidazol-1-yl]-3-[(cyclohexylmethyl)oxy]-2-thiophenecarbonitrile

1-[5-cyano-4-({[2-(trifluoromethyl)phenyl]methyl}oxy)-2-thienyl]-N-[2-(4-morpholinyl)ethyl]-1H-benzimidazole-5-carboxamide
5-[5,6-bis(methyloxy)-1H-benzimidazol-1-yl]-3-(((1R)-1-[2-(trifluoromethyl)phenyl]ethyl}oxy)-2-thiophenecarbonitrile
5-[5,6-bis(methyloxy)-1H-benzimidazol-1-yl]-3-[(2-phenylethyl)oxy]-2-thiophenecarbonitrile
5-[5,6-bis(methyloxy)-1H-benzimidazol-1-yl]-3-[(2-((trifluoromethyl)oxy]phenyl)methyl}oxy)-2-thiophenecarbonitrile
5-[5,6-bis(methyloxy)-1H-benzimidazol-1-yl]-3-({[2-(methyloxy)phenyl]methyl}oxy)-2-thiophenecarbonitrile
5-[5,6-bis(methyloxy)-1H-benzimidazol-1-yl]-3-[[2-(4-morpholinyl)ethyl]oxy]-2-thiophenecarbonitrile
5-[5,6-bis(methyloxy)-1H-benzimidazol-1-yl]-3-[[2-(2-oxo-1-pyrrolidinyl)ethyl]oxy]-2-thiophenecarbonitrile
5-[5,6-bis(methyloxy)-1H-benzimidazol-1-yl]-3-[(tetrahydro-2-furanylmethyl)oxy]-2-thiophenecarbonitrile
5-[5,6-bis(methyloxy)-1H-benzimidazol-1-yl]-3-[(tetrahydro-2H-pyran-2-ylmethyl)oxy]-2-thiophenecarbonitrile
5-[5,6-bis(methyloxy)-1H-benzimidazol-1-yl]-3-[[2-(phenyloxy)ethyl]oxy]-2-thiophenecarbonitrile
5-[5,6-bis(methyloxy)-1H-benzimidazol-1-yl]-3-[[1S)-1-(2-chlorophenyl)butyl]oxy]-2-thiophenecarbonitrile
5-[5,6-bis(methyloxy)-1H-benzimidazol-1-yl]-3-[(3-thienylmethyl)oxy]-2-thiophenecarbonitrile
5-[5,6-bis(methyloxy)-1H-benzimidazol-1-yl]-3-[(2-thienylmethyl)oxy]-2-thiophenecarbonitrile
5-[5,6-bis(methyloxy)-1H-benzimidazol-1-yl]-3-[[1R)-1-methylpropyl]oxy]-2-thiophenecarbonitrile

or a salt, solvate or physiologically functional derivative thereof.

5 8. Use of a compound of Formula (Ia), or a salt, solvate, or a physiologically functional derivative thereof in the preparation of a medicament for use in the treatment of a disorder mediated by inappropriate IKK3 activity.

10 9. Use according to claim 7 wherein disorder mediated by inappropriate IKK3 activity is inflammatory bowel disease, asthma and COPD (chronic obstructive pulmonary disease); osteoarthritis, osteoporosis and fibrotic diseases; dermatosis, including psoriasis, atopic dermatitis and ultraviolet radiation (UV)-induced skin damage; autoimmune diseases including systemic lupus erythematosus, multiple sclerosis, psoriatic arthritis, ankylosing spondylitis, tissue and organ rejection, Alzheimer's disease, stroke, atherosclerosis, restenosis, diabetes, glomerulonephritis, cancer, including Hodgkins

disease, cachexia, inflammation associated with infection and certain viral infections, including acquired immune deficiency syndrome (AIDS), adult respiratory distress syndrome, and Ataxia Telangiectasia.

- 5 10. Use according to claim 8 or 9 where the compound of Formula (Ia) is selected from:

5-[5,6-bis(methyloxy)-1 <i>H</i> -benzimidazol-1-yl]-3-({[4-(hydroxymethyl)phenyl]methyl}oxy)-2-thiophenecarbonitrile
5-[5,6-bis(methyloxy)-1 <i>H</i> -benzimidazol-1-yl]-3-({[(1 <i>R</i> )-1-(2-chlorophenyl)ethyl]oxy}-2-thiophenecarbonitrile
5-(1 <i>H</i> -benzimidazol-1-yl)-3-({[2-methylphenyl]methyl}oxy)-2-thiophenecarbonitrile
5-(1 <i>H</i> -benzimidazol-1-yl)-3-[(phenylmethyl)oxy]-2-thiophenecarbonitrile
5-[5,6-bis(methyloxy)-1 <i>H</i> -benzimidazol-1-yl]-3-({[(1-methyl-1 <i>H</i> -1,2,3-benzotriazol-6-yl)methyl]oxy}-2-thiophenecarbonitrile)
5-(6-(methyloxy)-5-({[2-(4-morpholinyl)ethyl]oxy}-1 <i>H</i> -benzimidazol-1-yl)-3-({[2-(trifluoromethyl)phenyl]methyl}oxy)-2-thiophenecarbonitrile
5-[5,6-bis(methyloxy)-1 <i>H</i> -benzimidazol-1-yl]-3-({[(2,5-difluorophenyl)methyl]oxy}-2-thiophenecarbonitrile
5-[5,6-bis(methyloxy)-1 <i>H</i> -benzimidazol-1-yl]-3-({[2-(trifluoromethyl)phenyl]methyl}oxy)-2-thiophenecarbonitrile
5-[5,6-bis(methyloxy)-1 <i>H</i> -benzimidazol-1-yl]-3-({[2-(methylsulfonyl)phenyl]methyl}oxy)-2-thiophenecarbonitrile
5-(5-chloro-1 <i>H</i> -benzimidazol-1-yl)-3-[(phenylmethyl)oxy]-2-thiophenecarbonitrile
5-[5,6-bis(methyloxy)-1 <i>H</i> -benzimidazol-1-yl]-3-({[(2,6-difluorophenyl)methyl]oxy}-2-thiophenecarbonitrile
5-[5-({[3-hydroxypropyl]oxy}-6-(methyloxy)-1 <i>H</i> -benzimidazol-1-yl]-3-({[2-(trifluoromethyl)phenyl]methyl}oxy)-2-thiophene-carbonitrile
5-[5,6-bis(methyloxy)-1 <i>H</i> -benzimidazol-1-yl]-3-({[(3-bromo-4-pyridinyl)methyl]oxy}-2-thiophenecarbonitrile
5-[5,6-bis(methyloxy)-1 <i>H</i> -benzimidazol-1-yl]-3-[(cyclohexylmethyl)oxy]-2-thiophenecarbonitrile
1-[5-cyano-4-({[2-(trifluoromethyl)phenyl]methyl}oxy)-2-thienyl]- <i>N</i> -[2-(4-morpholinyl)ethyl]-1 <i>H</i> -benzimidazole-5-carboxamide
5-[5,6-bis(methyloxy)-1 <i>H</i> -benzimidazol-1-yl]-3-({[(1 <i>R</i> )-1-[2-(trifluoromethyl)phenyl]ethyl]oxy}-2-thiophenecarbonitrile
5-[5,6-bis(methyloxy)-1 <i>H</i> -benzimidazol-1-yl]-3-[(2-phenylethyl)oxy]-2-thiophenecarbonitrile
5-[5,6-bis(methyloxy)-1 <i>H</i> -benzimidazol-1-yl]-3-({[2-[(trifluoromethyl)oxy]phenyl]methyl}oxy)-2-thiophenecarbonitrile
5-[5,6-bis(methyloxy)-1 <i>H</i> -benzimidazol-1-yl]-3-({[2-(methyloxy)phenyl]methyl}oxy)-2-thiophenecarbonitrile

5-[5,6-bis(methyloxy)-1 <i>H</i> -benzimidazol-1-yl]-3-[[2-(4-morpholinyl)ethyl]oxy]-2-thiophenecarbonitrile
5-[5,6-bis(methyloxy)-1 <i>H</i> -benzimidazol-1-yl]-3-[[2-(2-oxo-1-pyrrolidinyl)ethyl]oxy]-2-thiophenecarbonitrile
5-[5,6-bis(methyloxy)-1 <i>H</i> -benzimidazol-1-yl]-3-[(tetrahydro-2-furanylmethyl)oxy]-2-thiophenecarbonitrile
5-[5,6-bis(methyloxy)-1 <i>H</i> -benzimidazol-1-yl]-3-[(tetrahydro-2 <i>H</i> -pyran-2-ylmethyl)oxy]-2-thiophenecarbonitrile
5-[5,6-bis(methyloxy)-1 <i>H</i> -benzimidazol-1-yl]-3-[[2-(phenyloxy)ethyl]oxy]-2-thiophenecarbonitrile
5-[5,6-bis(methyloxy)-1 <i>H</i> -benzimidazol-1-yl]-3-[[[(1 <i>S</i> )-1-(2-chlorophenyl)butyl]oxy]-2-thiophenecarbonitrile
5-[5,6-bis(methyloxy)-1 <i>H</i> -benzimidazol-1-yl]-3-[(3-thienylmethyl)oxy]-2-thiophenecarbonitrile
5-[5,6-bis(methyloxy)-1 <i>H</i> -benzimidazol-1-yl]-3-[(2-thienylmethyl)oxy]-2-thiophenecarbonitrile
5-[5,6-bis(methyloxy)-1 <i>H</i> -benzimidazol-1-yl]-3-[[[(1 <i>R</i> )-1-methylpropyl]oxy]-2-thiophenecarbonitrile

or a salt, solvate or physiologically functional derivative thereof.

